

Customer No. 20350

TOWNSEND and TOWNSEND and CREW LLP
Two Embarcadero Center, 8th Floor
San Francisco, California 94111-3834
(415) 576-0200

**ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
Washington, D.C. 20231**

11/28/00



JC957 U.S. PTO

Attorney Docket No. 17682A-003610US

Client Ref No. 152A

"Express Mail" Label No. EL623997682US

Date of Deposit: November 28, 2000



Sir:

Transmitted herewith for filing under 37 CFR 1.53(b) is the

- ☐ patent application of
- ☐ continuation patent application of
- ☐ divisional patent application of
- ☒ continuation-in-part patent application of

Inventor(s)/Applicant Identifier: **Robert O. Dempcy, Alexander A. Gall, Sergey G. Lokhov, Irina A. Afonina, Michael J. Singer, Igor V. Kutyavin, Nicolaas M.J. Vermeulen**

For: **MODIFIED OLIGONUCLEOTIDES FOR MISMATCH DISCRIMINATION**

- ☒ This application claims priority from each of the following Application Nos./filing dates:
This application Application No. 09/640,953 filed August 16, 2000, 09/054,832, filed April 3, 1998; 09/431,385, filed November 1, 1999, 09/054,830, filed April 3, 1998, now U.S. Patent No. 6,127,121, the disclosure(s) of which is (are) incorporated by reference.
- ☐ Please amend this application by adding the following before the first sentence: This application is a continuation-in-part of U.S. Application, the disclosures of each being incorporated herein by reference.

Enclosed are:


- ☒ Title Page
- ☒ 87 page(s) of specification
- ☒ 18 page(s) of claims
- ☒ 1 page of Abstract
- ☒ 5 sheet(s) of ☐ formal ☒ informal drawing(s).
- ☒ A ☐ signed ☒ unsigned Declaration.
- ☐

**In view of the Unsigned Declaration as filed with this application and pursuant to 37 CFR §1.53(f),
Applicant requests deferral of the filing fee until submission of the Missing Parts of Application.**

DO NOT CHARGE THE FILING FEE AT THIS TIME.

Telephone:
(415) 576-0200

Facsimile:
(415) 576-0300


William B. Kezer
Reg No.: 37,369
Attorneys for Applicant

PATENT APPLICATION

Modified Oligonucleotides for Mismatch Discrimination

Inventor(s):

Robert O. Dempcy, a citizen of the United States
residing at 11421 NE 115th Court, Kirkland, Washington 98033

Alexander A. Gall, a citizen of Russia
residing at 20051 170th Avenue, NE, Woodinville, Washington 98072

Sergey G. Lokhov, a citizen of Russia
residing at 13215 NE, 123rd Street, #312, Kirkland, Washington 98034

Irina A. Afonina, a citizen of the United States
residing at 2717 144th Court SE, Mill Creek, Washington 98012

Michael J. Singer, a citizen of the United States
residing at 11526 36th Avenue, NE, Seattle, Washington 98125

Igor V. Kutyavin, a citizen of Russia
residing at 18943 203rd Avenue, NE, Woodinville, Washington 98072

Nicolaas M.J. Vermeulen, a citizen of the United States
residing at 19334 196th Avenue, NE, Woodinville, Washington 98072

Assignee:

Epoch Biosciences, Inc.
21720 – 23rd Dr., SE, #150
Bothell, WA 98021

Entity: Small

AS FILED NOVEMBER 28, 2000

Modified Oligonucleotides for Mismatch Discrimination

CROSS-REFERENCES TO RELATED APPLICATIONS

5 This application is a continuation-in-part of U.S. Application No. 09/640,953
filed August 16, 2000, which is a continuation of U.S. Application Ser. No. 09/054,832, filed
April 3, 1998; and a continuation-in-part of U.S. Application Ser. No. 09/431,385, filed
November 1, 1999, which is a continuation of U.S. Application Ser. No. 09/054,830, filed
April 3, 1998, now U.S. Patent No. 6,127,121; and claims the benefit of Provisional
10 application Ser. No. 60/186,046, filed March 1, 2000, the disclosures of each being
incorporated herein by reference.

STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

15 Not applicable.

BACKGROUND OF THE INVENTION

20 This application is in the field of molecular biology relating to the use of
oligonucleotides as probes and primers in liquid, solid and mixed phase assays. The
application further relates to the use of modified nucleic acid bases and modified
oligonucleotides to improve the hybridization properties and discriminatory abilities of
oligonucleotides that are used in arrays and as probes and primers.

25 Many techniques currently in use in molecular biology utilize oligonucleotides
as probes and/or primers. It is often advantageous, in the practice of these techniques, to be
able to distinguish between two or more sequences which are related but which differ by one
or more nucleotides. For example, many mutations of clinical significance differ by only a
single nucleotide from the wild-type sequence. Polymorphisms in mammalian genomes are
also often characterized by sequence differences of one or a few nucleotides. The ability to
30 make such a distinction is known as mismatch discrimination. In practical terms, mismatch
discrimination describes the property by which a defined sequence oligonucleotide, at a given
stringency, hybridizes strongly (one manifestation of which is that the hybrids have a high
melting temperature) to a target sequence with which it is complementary along its entire
length (a perfect hybrid or perfect match), but hybridizes detectably more weakly to a target

sequence that is non-complementary to the sequence of the oligonucleotide at one or a few nucleotides (a mismatch). The differences in hybridization strength are such that a particular stringency can be selected at which a perfect match is detectable as a hybrid and a mismatch fails to form a hybrid.

5 In a nucleic acid duplex, each base pair contributes to stability. Hence, the shorter the duplex, the greater the relative contribution of each individual base pair to the stability of the duplex. As a result, the difference in stability between a perfect match and a mismatch will be greater for shorter oligonucleotides. However, short oligonucleotides hybridize weakly, even to a perfectly complementary sequence, and thus must be hybridized
10 under conditions of reduced stringency. Thus, the potential discriminatory power of short oligonucleotides cannot be easily realized except under conditions of low stringency.

What is needed in the art are new methods for mismatch discrimination, particularly for single-nucleotide mismatches, under conditions of high stringency; for example, at the elevated temperatures characteristic of most nucleic acid amplification
15 reactions. Surprisingly, the present invention provides such methods, along with new reagents and compositions which can be used in the methods.

SUMMARY OF THE INVENTION

20 The present invention provides a number of modified oligonucleotides found to have exceptional properties and usefulness in a variety of assays. Accordingly, the present invention also provides methods for using the modified oligonucleotides described herein.

In one aspect, the present invention provides modified oligonucleotides having at least two bases selected from unsubstituted and 3-substituted pyrazolo[3,4-d]pyrimidine
25 bases. In preferred embodiments, the oligonucleotides having modified bases will further comprise other moieties such as detectable labels, fluorescence and chemiluminescence quenchers and/or minor groove binders and/or other types of modified bases or base analogs.

In another aspect, the present invention provides modified oligonucleotides having at least one 5-substituted pyrimidine base and at least one 3-substituted pyrazolo[3,4-
30 d]pyrimidine base. In preferred embodiments, these modified oligonucleotides will further comprise other moieties (as above) such as detectable labels, fluorescence and chemiluminescence quenchers and/or minor groove binders.

In yet another aspect, the present invention provides methods for distinguishing polynucleotides having related sequences.

In still another aspect, the present invention provides methods for detecting the presence of a target sequence in a polynucleotide.

In still other aspects, the present invention provides methods for primer extension, and methods for determining the nucleotide sequence of a polynucleotide.

5 In related aspects, the present invention provides methods for examining gene expression in a cell, and methods for identifying a mutation or polymorphism in a target sequence of a gene of interest.

In still another aspect, the present invention provides a number of modified bases that are useful in preparing modified oligonucleotides for the methods described herein
10 and other conventional assays and methods.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides structures for several modified bases and their
15 abbreviations. The wavy line is used to denote the position of an attached sugar moiety (unprotected, protected, activated and the like).

Figure 2 is a graph illustrating the balancing of the T_m of 8-mer GC-rich and AT-rich probes with different combinations of MGB, PPPA and PU.

Figure 3 is a chart which illustrates one advantage achieved through the use of
20 PPPA and PPG in MGB-modified oligonucleotide probes. As seen in the figure, the modified bases allow shortening of the probe that shows increased mismatch discrimination in real-time PCR. \tilde{A} is PPPA and \tilde{G} is PPG.

Figure 4 illustrates an InvaderTM assay in which the modified oligonucleotides of the invention can be used.

25 **Figure 5** illustrates a comparison of InvaderTM probe performance with different numbers of PPGs.

DESCRIPTION OF THE INVENTION

Abbreviations and Definitions

30 Abbreviations for a number of the modified bases described herein are provided as follows (structures of these bases are shown in Figure 1): 6-amino-3-prop-1-ynyl-

The term "alkynyl" refers to a linear monovalent hydrocarbon radical or a branched monovalent hydrocarbon radical containing at least one triple bond and having the number of carbon atoms indicated in the prefix. For example, (C₂-C₆)alkynyl is meant to include ethynyl, propynyl, and the like.

The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively. Similarly, the term dialkylamino refers to an amino group having two attached alkyl groups that can be the same or different.

The term "aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms which is unsubstituted or substituted independently with one to four substituents, preferably one, two, or three substituents selected from alkyl, cycloalkyl, cycloalkyl-alkyl, halo, nitro, cyano, hydroxy, alkoxy, amino, acylamino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, heteroalkyl, COR (where R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), -(CR'R'')_n-COOR (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl) or -(CR'R'')_n-CONR^aR^b (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R^a and R^b are, independently of each other, hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, phenyl or phenylalkyl). More specifically the term aryl includes, but is not limited to, phenyl, biphenyl, 1-naphthyl, and 2-naphthyl, and the substituted forms thereof. Similarly, the term "heteroaryl" refers to those aryl groups wherein one or more heteroatoms or heteroatom functional groups have replaced a ring carbon, while retaining aromatic properties, e.g., pyridyl, quinoliny, quinazolinyl, thienyl, and the like. For brevity, the term aryl, when used in combination with other radicals (e.g., aryloxy, arylalkyl) is meant to include both aryl groups and heteroaryl groups as described above.

The term "arylalkyl" refers to a radical -R^aR^b where R^a is an alkylene group (having the indicated number of carbon atoms, or if unspecified having six or fewer main chain carbon atoms) and R^b is an aryl group as defined herein. Examples of arylalkyl groups include benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like.

Similarly the term "arylalkenyl" means a radical $-R^aR^b$ where R^a is an alkenylene group and R^b is an aryl group as defined herein, *e.g.*, 3-phenyl-2-propenyl, and the like.

"Arylheteroalkyl" means a radical $-R^aR^b$ where R^a is an heteroalkylene group (having the indicated number of carbon atoms) and R^b is an aryl group as defined herein, *e.g.*, 2-hydroxy-2-phenyl-ethyl, 2-hydroxy-1-hydroxymethyl-2-phenyl-ethyl, and the like.

The term "aryloxy", refers to a radical $-OR$ where R is an aryl group, *e.g.*, phenoxy, naphthyloxy and the like.

The prefix "halo" and the term "halogen" when used to describe a substituent, refer to $-F$, $-Cl$, $-Br$ and $-I$.

The term "heteroalkyl" refers to an alkyl radical as defined herein with one, two or three substituents independently selected from cyano, $-OR^a$, $-NR^bR^c$, and $-S(O)_nR^d$ (where n is an integer from 0 to 2), with the understanding that the point of attachment of the heteroalkyl radical is through a carbon atom of the heteroalkyl radical. R^a is hydrogen, alkyl, aryl, arylalkyl, alkoxycarbonyl, aryloxycarbonyl, carboxamido, or mono- or di-alkylcarbamoyl. R^b is hydrogen, alkyl, aryl or arylalkyl. R^c is hydrogen, alkyl, aryl, arylalkyl, alkoxycarbonyl, aryloxycarbonyl, carboxamido, mono- or di-alkylcarbamoyl or alkylsulfonyl. R^d is hydrogen (provided that n is 0), alkyl, aryl, arylalkyl, amino, mono-alkylamino, di-alkylamino, or hydroxyalkyl. Representative examples include, for example, 2-hydroxyethyl, 2,3-dihydroxypropyl, 2-methoxyethyl, benzyloxymethyl, 2-cyanoethyl, and 2-methylsulfonyl-ethyl. For each of the above, R^a , R^b , R^c , and R^d can be further substituted by NH_2 , fluorine, alkylamino, di-alkylamino, OH or alkoxy. Additionally, the prefix indicating the number of carbon atoms (*e.g.*, C_1 - C_{10}) refers to the total number of carbon atoms in the portion of the heteroalkyl group exclusive of the cyano, $-OR^a$, $-NR^bR^c$, or $-S(O)_nR^d$ portions.

The term "heterocyclyl" refers to a saturated or unsaturated non-aromatic cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from O , NR (where R is independently hydrogen or alkyl) or $S(O)_n$ (where n is an integer from 0 to 2), the remaining ring atoms being C , where one or two C atoms may optionally be replaced by a carbonyl group. The heterocyclyl ring may be optionally substituted independently with one, two, or three substituents selected from alkyl, halo, nitro, cyano, hydroxy, alkoxy, amino, mono-alkylamino, di-alkylamino, haloalkyl, haloalkoxy, $-COR$

(where R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), $-(CR'R'')_n$ -COOR (n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl, and R is hydrogen, alkyl, cycloalkyl, cycloalkyl-alkyl, phenyl or phenylalkyl), or $-(CR'R'')_n$ -

CONR^aR^b (where n is an integer from 0 to 5, R' and R'' are independently hydrogen or alkyl,

5 and R^a and R^b are, independently of each other, hydrogen, alkyl, phenyl or phenylalkyl).

More specifically the term heterocyclyl includes, but is not limited to, tetrahydropyranyl, piperidino, N-methylpiperidin-3-yl, piperazino, N-methylpyrrolidin-3-yl, 3-pyrrolidino, 2-pyrrolidon-1-yl, morpholino, thiomorpholino, thiomorpholino-1-oxide, thiomorpholino-1,1-dioxide, pyrrolidinyl, and the derivatives thereof. The prefix indicating the number of carbon
10 atoms (e.g., C₃-C₁₀) refers to the total number of carbon atoms in the portion of the heterocyclyl group exclusive of the number of heteroatoms.

The terms "heterocyclylalkyl," "heterocyclylalkenyl," "heterocyclylalkynyl" refer to radicals -R^aR^b where R^a is an alkylene, alkenylene or alkynylene group, respectively, and R^b is a heterocyclyl group as defined herein, e.g., tetrahydropyran-2-ylmethyl, 4-methylpiperazin-1-ylethyl, 3-piperidinylmethyl, and the like.
15

The terms "heteroalkylene" means a linear saturated divalent hydrocarbon radical of one to six carbons or a branched saturated hydrocarbon radical of three to six carbon atoms with one, two or three substituents independently selected from -OR^a, -NR^bR^c, and -S(O)_nR^d (where n is an integer from 0 to 2) where, R^a, R^b, R^c, and R^d are as defined
20 herein for a heteroalkyl radical. Examples include, 2-hydroxyethan-1,2-diyl, 2-hydroxypropan-1,3-diyl and the like.

Each of the above terms (e.g., "alkyl," "heteroalkyl," and "aryl") are meant to include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

25 Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, heterocycloalkyl, and heterocycloalkenyl) can be a variety of groups selected from: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R'', -NR''C(O)₂R', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -CN
30 and -NO₂ in a number ranging from zero to four, preferably, zero, one, two or three substituents. R', R'' and R''' each independently refer to hydrogen, unsubstituted (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl, aryl substituted with 1-3 halogens, unsubstituted

alkyl, alkoxy or thioalkoxy groups, or aryl-(C₁-C₄)alkyl groups. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will understand that the term "alkyl" in its broadest sense is meant to include groups such as haloalkyl (e.g., -CF₃ and -CH₂CF₃) and acyl (e.g., -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like). Preferably, the alkyl groups will have from 0-3 substituents, more preferably 0, 1, or 2 substituents, unless otherwise specified.

Similarly, substituents for the aryl groups are varied and are selected from: - halogen, -OR', -OC(O)R', -NR'R'', -SR', -R', -CN, -NO₂, -CO₂R', -CONR'R'', -C(O)R', -OC(O)NR'R'', -NR''C(O)R', -NR''C(O)₂R', -NR'-C(O)NR''R'', -NH-C(NH₂)=NH, -NR'C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -N₃, -CH(Ph)₂, perfluoro(C₁-C₄)alkoxy, and perfluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'' and R''' are independently selected from hydrogen, (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl.

Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CH₂)_q-U-, wherein T and U are independently -NH-, -O-, -CH₂- or a single bond, and q is an integer of from 0 to 2.

Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CH₂-, -O-, -NH-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 3. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -(CH₂)_s-X-(CH₂)_t-, where s and t are independently integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituent R' in -NR'- and -S(O)₂NR'- is selected from hydrogen or unsubstituted (C₁-C₆)alkyl.

Certain compounds or oligonucleotides of the present invention may exist in a salt form. Such salts include base addition salts such as sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When the compounds or modified oligonucleotides of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of

acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from organic acids like acetic, propionic, isobutyric, maleic, malonic, lactic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are all intended to be encompassed within the scope of the present invention. The methods for the determination of stereochemistry and the separation of isomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 1992).

The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the

compounds of the present invention, whether radioactive or not (e.g., ^2H), are intended to be encompassed within the scope of the present invention.

"Protecting group" or "protected form thereof" refers to a grouping of atoms that when attached to a reactive group in a molecule masks, reduces or prevents that reactivity. Examples of protecting groups can be found in T.W. Greene and P.G. Futs, *Protective Groups in Organic Chemistry*, (Wiley, 2nd ed. 1991) and Harrison and Harrison *et al.*, *Compendium of Synthetic Organic Methods*, Vols. 1-8 (John Wiley and Sons. 1971-1996). Representative amino protecting groups include formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethyl silyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (FMOC), nitro-veratryloxycarbonyl (NVOC) and the like. Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as benzyl and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers and allyl ethers.

"Optional" or "optionally" in the above definitions means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "heterocyclo group optionally mono- or di- substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the heterocyclo group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques in organic chemistry, biochemistry, oligonucleotide synthesis and modification, bioconjugate chemistry, nucleic acid hybridization, molecular biology, microbiology, genetics, recombinant DNA, and related fields as are within the skill of the art. These techniques are fully explained in the literature. See, for example, Maniatis, Fritsch & Sambrook, *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory Press (1982); Sambrook, Fritsch & Maniatis, *MOLECULAR CLONING: A LABORATORY MANUAL*, Second Edition, Cold Spring Harbor Laboratory Press (1989); Ausubel, *et al.*, *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John Wiley & Sons (1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996); Gait (ed.), *OLIGONUCLEOTIDE SYNTHESIS: A PRACTICAL APPROACH*, IRL Press (1984);

General

5 Oligonucleotides are short polymers of nucleotides, generally less than 200 nucleotides, preferably less than 150 nucleotides, more preferably less than 100 nucleotides, more preferably less than 50 nucleotides and most preferably less than 21 nucleotides in length. Polynucleotides are generally considered, in the art, to comprise longer polymers of nucleotides than do oligonucleotides, although there is an art-recognized overlap between the upper limit of oligonucleotide length and the lower limit of polynucleotide length. With respect to the present invention, "oligonucleotide" generally refers to a nucleic acid, usually comprising a detectable label, that is used as a probe or as a primer; while polynucleotide refers to a nucleic acid containing a target sequence. Consequently, for the purposes of the present invention, the terms "oligonucleotide" and "polynucleotide" shall not be considered
10 limiting with respect to polymer length.

 The present invention provides modified oligonucleotides having new and surprising properties of superior mismatch discrimination, compared to unmodified oligonucleotides. Modified oligonucleotides of the invention are used as probes, wherein their hybridization to a target sequence is detected, or as primers, wherein their hybridization to a target sequence is followed by polynucleotide synthesis initiated from the 3' terminus of the modified oligonucleotide, and the synthesized product (*i.e.*, the extension product) is detected.
20 detected.

 A target sequence refers to a nucleotide sequence which comprises a site of hybridization for a probe or a primer. Target sequences can be found in any nucleic acid including, but not limited to, genomic DNA, cDNA, RNA and any amplified product thereof, and can comprise a wild-type gene sequence, a mutant gene sequence, a non-coding sequence, a regulatory sequence, *etc.* A target sequence will generally be less than 100 nucleotides, preferably less than 50 nucleotides, and most preferably, less than 21 nucleotides in length.
25 in length.

Description of the Embodiments

5 The present invention provides a number of modified oligonucleotides which can generally be divided into three groups.

The first group of modified oligonucleotides are those having at least two modified bases replacing the naturally-occurring bases. Here, the modified bases will be unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines. In some embodiments, however, the modified bases will be selected such that at least one of the bases is a 5-substituted
10 pyrimidine and at least one of the bases is an unsubstituted or a 3-substituted pyrazolo[3,4-d]pyrimidine. Preferably, this group of modified oligonucleotides will have additional attached groups (e.g., minor groove binders, reporter groups, quenchers, etc.) which assist during assays to detect target sequences.


The second group of modified oligonucleotides are those having at least one
15 modified base, but also having an attached minor groove binder, reporter group, quencher, or the like.

The third group of modified oligonucleotides are those that comprise one or more of the novel modified bases described below. As in the first group, this group of modified oligonucleotides will preferably have attached groups selected from, for example,
20 minor groove binders, reporter groups or quenchers.

Modified Oligonucleotides


In one aspect, the present invention provides modified oligonucleotides
25 comprising at least two bases selected from the group consisting of unsubstituted and 3-substituted pyrazolo[3,4-d]pyrimidine bases.

The pyrazolo[3,4-d]pyrimidine bases are those bases in which a pyrazole ring is fused to a pyrimidine ring in the following specific orientation:



Ia


in which the wavy line indicates the point of attachment between the base and either an attached oligomer-forming sugar or an amino acid involved in peptide nucleic acid formation. Additionally, the groups X^1 and X^2 are independently H, OH or NH_2 such that the pyrazolo[3,4-d]pyrimidine bases approximate the construction of the natural purine bases, guanosine, adenine and inosine, as well as related derivatives of those bases. In this group of embodiments, an “unsubstituted pyrazolo[3,4-d]pyrimidine base” refers to those bases of the general formula Ia, while the term “3-substituted pyrazolo[3,4-d]pyrimidine base” refers to those bases having the formula Ib, in which R^1 is defined as described below.



Ib

In formula Ib, the symbols X^1 and X^2 independently represent H, OH, NH_2 or a protected form thereof. The symbol R^1 represents a member selected from (C_1 - C_{12})heteroalkyl, (C_2 - C_{12})heteroalkenyl, (C_2 - C_{12})heteroalkynyl, -O-(C_1 - C_{12})alkyl, -O-(C_2 - C_{12})alkenyl, -O-(C_2 - C_{12})alkynyl, -S-(C_1 - C_{12})alkyl, -S-(C_2 - C_{12})alkenyl, -S-(C_2 - C_{12})alkynyl, heterocyclyl(C_1 - C_{12})alkyl, heterocyclyl(C_2 - C_{12})alkenyl, heterocyclyl(C_2 - C_{12})alkynyl, aryl(C_1 - C_{12})alkyl, aryl(C_2 - C_{12})alkenyl, aryl(C_2 - C_{12})alkynyl, aryl, heterocyclyl, halogen, -CN, -CONH₂ and protected forms thereof. Preferred heteroalkyl, heteroalkenyl and heteroalkynyl groups are those that terminate (distal to the pyrazolo[3,4-d]pyrimidine ring system) in a heteroatom group (e.g., OH, NH_2 , SH and the like). For example, preferred heteroalkyl, heteroalkenyl and heteroalkynyl groups include 3-amino-1-propyl, 4-hydroxy-1-butyl, 3-amino-1-propyn-1-yl, 4-hydroxy-3-hydroxymethyl-1-butyne-1-yl. Other preferred R^1 groups include those which terminate in an aryl or heterocyclic group (e.g., heterocyclyl(C_1 - C_{12})alkyl, heterocyclyl(C_2 - C_{12})alkenyl, heterocyclyl(C_2 - C_{12})alkynyl, aryl(C_1 - C_{12})alkyl, aryl(C_2 - C_{12})alkenyl and aryl(C_2 - C_{12})alkynyl). Preferred aryl and heterocyclic groups are either substituted or unsubstituted phenyl, thienyl, thiazolyl, imidazolyl, furanyl, oxazolyl, pyridinyl, pyrrolyl, indolyl, benzimidazolyl, benzthiazolyl and benzoxazolyl.

In particularly preferred embodiments, the pyrazolo[3,4-d]pyrimidine base is selected from:



Still more preferably, the modified bases are selected from Ic, Id or Ie wherein each R¹ is a propynyl, hydroxypropynyl, aminopropynyl, or a substituted or unsubstituted phenyl, thienyl, thiazolyl, imidazolyl, furanyl, oxazolyl, pyridinyl, pyrrolyl, indolyl, benzimidazolyl, benzthiazolyl or benzoxazolyl group attached directly to the pyrazolo[3,4-d]pyrimidine ring system or attached to the ring system via a one to four carbon linking group which can be saturated (e.g., ethylene, propylene, butylene) or unsaturated (e.g., acetylenyl, propynylene, butynylene, propenylene, butenylene).

The modified oligonucleotides of the present invention have a backbone such as, for example, that found with the naturally-occurring oligonucleotides or peptide-nucleic acids (e.g., heterocyclic bases attached to oligomer-forming sugars or peptide-nucleic acid-forming amino acids). Alternative oligomeric backbones are known to those of skill in the art (referred to below as “modified” backbones or comprising backbone modifications). For the purpose of this invention a modified oligonucleotide is any oligomeric polymer, including the natural phosphate backbone and non-natural backbones, that contains natural and/or modified bases and retains the ability to hybridize specifically to a complementary nucleic acid to form a stable duplex.

In preferred embodiments, the modified oligonucleotides of the invention have a backbone of sugar or glycosidic moieties, preferably 2-deoxyribofuranosides wherein all internucleotide linkages are the naturally occurring phosphodiester linkages. In alternative embodiments however, the 2-deoxy-β-D-ribofuranose groups are replaced with other sugars, for example, β-D-ribofuranose. In addition, β-D-ribofuranose may be present wherein the 2-OH of the ribose moiety is alkylated with a C₁₋₆ alkyl group (2-(O-C₁₋₆ alkyl) ribose) or with a C₂₋₆ alkenyl group (2-(O-C₂₋₆ alkenyl) ribose), or is replaced by a fluoro group (2-fluororibose). Related oligomer-forming sugars useful in the present invention are those that are “locked”, i.e., contain a methylene bridge between C-4' and an oxygen atom at C-2'. Other sugar moieties compatible with hybridization of the oligonucleotide can also be used, and are known to those of skill in the art, including, but not limited to, α-D-arabinofuranosides, α-2'-deoxyribofuranosides or 2',3'-dideoxy-3'-aminoribofuranosides. Oligonucleotides containing α-D-arabinofuranosides can be prepared as described in U.S.

Patent No. 5,177,196. Oligonucleotides containing 2',3'-dideoxy-3'-aminoribofuranosides are described in Chen et al. ((1995) *Nucleic Acids Res.* **23**:2661-2668). Synthetic procedures for locked nucleic acids (Singh et al, *Chem. Comm.*, 455-456 (1998); Wengel J., *Acc. Chem. Res.*, **32**:301-310 (1998)) and oligonucleotides containing 2'-halogen-2'-

deoxyribofuranosides (Palissa et al., *Z. Chem.* **27**:216 (1987)) have been described.. The phosphate backbone of the modified oligonucleotides described herein can also be modified so that the oligonucleotides contain phosphorothioate linkages and/or methylphosphonates and/or phosphoroamidates (Chen et al., *Nucl. Acids Res.*, **23**:2662-2668 (1995)).

Combinations of oligonucleotide linkages are also within the scope of the present invention.

Still other backbone modifications are known to those of skill in the art.

In another group of embodiments, the modified bases described herein are incorporated into PNA and DNA/PNA chimeras to balance T_m s and provide modified oligonucleotides having improved mismatch discrimination. Various modified forms of DNA and DNA analogues have been used in attempts to overcome some of the disadvantages of the use of DNA molecules as probes and primers. Among these are peptide nucleic acids (PNAs, also known as polyamide nucleic acids). Nielsen et al. (1991) *Science* **254**:1497-1500. PNAs contain heterocyclic base units, as found in DNA and RNA, that are linked by a polyamide backbone, instead of the sugar-phosphate backbone characteristic of DNA and RNA. PNAs are capable of hybridization to complementary DNA and RNA target sequences and, in fact, hybridize more strongly than a corresponding nucleic acid probe. The synthesis of PNA oligomers and reactive monomers used in the synthesis of PNA oligomers have been described in U.S. Patents No. 5,539,082; 5,714,331; 5,773,571; 5,736,336 and 5,766,855. Alternate approaches to PNA and DNA/PNA chimera synthesis and monomers for PNA synthesis have been summarized. Uhlmann et al. (1998) *Angew. Chem. Int. Ed.* **37**:2796-2823. Accordingly, the use of any combination of normal bases, unsubstituted pyrazolo[3,4-d]pyrimidine bases (e.g., PPG and PPA), 3-substituted pyrazolo[3,4-d]pyrimidines, modified purine, modified pyrimidine, 5-substituted pyrimidines, universal bases or a minor groove binder to balance the T_m of a PNA or DNA/PNA chimera is in the scope of this invention. The synthetic methods necessary for the synthesis of modified base monomeric units required for PNA and PNA/DNA chimeras synthesis are available in the art, see methods in this application and Uhlmann et al. *Angew. Chem. Int. Ed.* **37**:2796-2823 (1998).

Similarly, the present invention demonstrates a combination of normal bases, unsubstituted pyrazolo[3,4-d]pyrimidine bases (e.g., PPG and PPA), 3-substituted pyrazolo[3,4-d]pyrimidines, modified purine, modified pyrimidine, 5-substituted

pyrimidines, universal bases or a minor groove binder can be used to balance the T_m of any oligonucleotide polymer or oligopolymer/DNA chimera.

For the uses described herein, the modified oligonucleotides will preferably have from 4 to 70 bases, more preferably from 4 to 25 bases. In some embodiments, the modified oligonucleotides will have 15 or fewer, or more preferably 10 or fewer bases. Additionally, the modified oligonucleotides will, in some embodiments, have 3, 4, 5, 6, 7, 8, or 9 or more modified bases (either unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines). For each of the embodiments wherein multiple 3-substituted pyrazolo[3,4-d]pyrimidine bases are present, the modified bases can either be the same or different from each other.

In addition to the modified base components, the modified oligonucleotides of the present invention will in some embodiments comprise additional pendant groups such as, for example, intercalators, lipophilic groups, minor groove binders, reporter groups, chelating agents, quenchers and cross-linking agents attached to one or more of the internally located nucleotide bases, to the 3', to the 5' end, to both ends, or can have such pendant groups attached both internally and at one or both ends. Methods suitable for attaching intercalator, lipophilic groups, minor groove binders, reporter groups, chelating agents, quenchers and cross-linking agents to oligonucleotides have been described in, for example, U.S. Patent Nos. 5,512,667 and 5,419,966, PCT publication WO 96/32496 and U.S. application Ser. No. 09/457,616. The oligonucleotides of the invention can also have a relatively low molecular weight "tail moiety" attached either at the 3' or 5' end, or at both ends. By way of example a tail molecule can be a phosphate, a phosphate ester, an alkyl group, an aminoalkyl group, or a lipophilic group. The tail moiety can also link the intercalators, lipophilic groups, minor groove binders, reporter groups, chelating agents and cross-linking functionalities to the oligonucleotides of the invention.

The nature of tail moieties and methods for obtaining oligonucleotides with various tail moieties are also described in the above-referenced U.S. Patent Nos. 5,512,667 and 5,419,966.

Minor Groove Binders

In one group of embodiments, the modified oligonucleotide will also have a covalently attached minor groove binder (MGB). A variety of suitable minor groove binders have been described in the literature. See, for example, Kutyavin, et al. U.S. Patent No. 5,801,155; Wemmer, D.E., and Dervan P.B., *Current Opinion in Structural Biology*, 7:355-

361 (1997); Walker, W.L., Kopka, J.L. and Goodsell, D.S., *Biopolymers*, **44**:323-334 (1997); Zimmer, C & Wahnert, U. *Prog. Biophys. Molec. Bio.* **47**:31-112 (1986) and Reddy, B.S.P., Dondhi, S.M., and Lown, J. W., *Pharmacol. Therap.*, **84**:1-111 (1999).


Suitable methods for attaching MGBs (as well as reporter groups such as fluorophores and quenchers described below) through linkers to oligonucleotides are described in, for example, U.S. Patent Nos. 5,512,677; 5,419,966; 5,696,251; 5,585,481; 5,942,610 and 5,736,626.

The MGB can be attached at either or both ends of the oligonucleotide. In addition or alternatively, one or more MGBs can be attached in the interior of the oligonucleotide, depending on the length of the oligonucleotide. In general, conjugation of a MGB to either end of an oligonucleotide would provide the greatest degree of hybrid stability, since melting of an oligonucleotide duplex begins at the termini. Nonetheless, if both ends of a duplex formed by an oligonucleotide are relatively stable, for example, due to a high G+C content, attachment of a MGB in the interior of an oligonucleotide (for instance, near an A+T-rich sequence) could also enhance stability. The intended use of the MGB-oligonucleotide conjugate may also place limitations on the location of the conjugated MGB. For example, if an oligonucleotide is designed to be used as a primer, the 3'-hydroxy group must be free and capable of being elongated by a polymerizing enzyme. Alternatively, an assay that requires an oligonucleotide possessing a labeled 5'-end would require internal or 3'-end attachment of a MGB.

The location of a MGB within a MGB-modified oligonucleotide conjugate can also affect the discriminatory properties of such a conjugate. An unpaired region within a duplex will result in changes in the shape of the minor groove in the vicinity of the mispaired base(s). Since MGBs fit best within the minor groove of a perfectly-matched DNA duplex, mismatches resulting in shape changes in the minor groove would reduce binding strength of a MGB to a region containing a mismatch. Hence, the ability of a MGB to stabilize such a hybrid would be decreased, thereby increasing the ability of a MGB-oligonucleotide conjugate to discriminate a mismatch from a perfectly-matched duplex. On the other hand, if a mismatch lies outside of the region complementary to a MGB-oligonucleotide conjugate, discriminatory ability for unconjugated and MGB-conjugated oligonucleotides of equal length is expected to be approximately the same. Since the ability of an oligonucleotide probe to discriminate single base pair mismatches depends on its length, shorter oligonucleotides are more effective in discriminating mismatches. The primary advantage of the use of MGB-oligonucleotides conjugates in this context lies in the fact that much shorter

oligonucleotides compared to those used in the prior art (*i.e.*, 20-mers or shorter), having greater discriminatory powers, can be used, due to the pronounced stabilizing effect of MGB conjugation.

Preferred minor groove binders are those selected from the formulae:



the subscript *m* is an integer of from 2 to 5; the subscript *r* is an integer of from 2 to 10; and each *R^a* and *R^b* is independently a linking group to the modified oligonucleotide, H, -OR^c, -NR^cR^d, -COOR^c or -CONR^cR^d, wherein each *R^c* and *R^d* is selected from H, (C₁-C₁₂)heteroalkyl, (C₂-C₁₂)heteroalkenyl, (C₂-C₁₂)heteroalkynyl, (C₁-C₁₂)alkyl, (C₂-C₁₂)alkenyl, (C₂-C₁₂)alkynyl, aryl(C₁-C₁₂)alkyl and aryl.

Particularly preferred minor groove binders include the trimer of 3-carbamoyl-1,2-dihydro-(3*H*)-pyrrolo[3,2-*e*]indole-7-carboxylate (CDPI₃), the pentamer of N-methylpyrrole-4-carbox-2-amide (MPC₅) and other minor groove binders that exhibit increased mismatch discrimination. Additional MGB moieties that will find use in the practice of the present invention are disclosed in co-owned U.S. Patent No. 5,801,155. In certain embodiments, the MGBs can have attached water solubility-enhancing groups (*e.g.*, sugars or amino acids).

Reporter Groups

In another group of embodiments, the modified oligonucleotide will further comprise at least one covalently attached reporter group. The reporter groups can be attached using methods and linking groups described above for the MGBs. Suitable reporter groups for the present modified oligonucleotides include beads, nanoparticles (Taton, T.A. et al, *Science* **289**:1757-1760 (2000)), chemiluminescers, isotopes, enzymes and fluorophores. Preferably, the reporter group is a fluorophore (see, Haugland, R.P., HANDBOOK OF FLUORESCENT PROBES AND RESEARCH CHEMICALS, Sixth Edition, Molecular Probes, Eugene, OR, 1996). Suitable fluorophores include the resorufin dyes, coumarin dyes, rhodamine dyes, cyanine dyes, BODIPY dyes and pyrenes.

Quenchers

Recently developed detection methods employ the process of fluorescence resonance energy transfer (FRET) for the detection of probe hybridization rather than direct detection of fluorescence intensity. In this type of assay, FRET occurs between a donor fluorophore (reporter) and an acceptor molecule (quencher) when the absorption spectrum of the quencher molecule overlaps with the emission spectrum of the donor fluorophore and the two molecules are in close proximity. The excited-state energy of the donor fluorophore is transferred to the neighboring acceptor by a resonance dipole-induced dipole interaction, which results in quenching of the donor fluorescence. If the acceptor molecule is a fluorophore, its fluorescence may sometimes be increased. The efficiency of the energy transfer between the donor and acceptor molecules is highly dependent on distance between the molecules. Equations describing this relationship are known. The Forster distance (R_0) is described as the distance between the donor and acceptor molecules where the energy transfer is 50% efficient. Other mechanisms of fluorescence quenching are also known, such as, collisional and charge transfer quenching. There is extensive guidance in the art for selecting quencher and fluor pairs and their attachment to oligonucleotides (Haugland, R.P., HANDBOOK OF FLUORESCENT PROBES AND RESEARCH CHEMICALS, Sixth Edition, Molecular Probes, Eugene, OR, 1996; U.S. Patent Nos. 3,996,345 and 4,351,760 and the like). Preferred quenchers are described in co-owned U.S. Ser. No. 09/457,616 (filed December 8, 1999).

Fluorophores and Quenchers

In certain embodiments of the present invention, oligonucleotides comprising fluorescent labels (fluorophores) and/or fluorescence quenching agents are used. In a preferred embodiment, an oligonucleotide contains both a fluorophore and a quenching agent. Fluorescent labels include, but are not limited to, fluoresceins, rhodamines, cyanines, phycoerythrins, and other fluorophores as described herein. Still other suitable fluorophores are known to those of skill in the art. As noted above, quenching agents or quenchers can absorb energy emitted by a fluorophore so as to reduce the amount of fluorescence emitted (*i.e.*, quench the emission of the fluorescent label). Different fluorophores are quenched by different quenching agents. In general, the spectral properties of a particular fluorophore/quenching agent pair are such that one or more absorption wavelengths of the


quencher overlaps one or more of the emission wavelengths of the fluorophore. A preferred fluorophore/quencher pair can be selected by one of skill in the art by comparison of emission and excitation wavelengths according to the properties set forth above.

For use in amplification assays conducted at elevated temperatures, such as a polymerase chain reaction, or other procedures utilizing thermostable enzymes, the label is preferably one that is stable at elevated temperatures. For assays involving polymerization, the label is one that does not interfere with the activity of the polymerizing enzyme.

Additionally, the label will be present at the 5' and/or 3' end of the oligonucleotide, and/or can also be present internally in a non-interfering position. Accordingly, the label can be attached to any of the bases, sugars or phosphate moieties of the oligonucleotide, or to any linking group that is itself attached to one of these moieties.

While the invention has been described in terms of the various bases and other optional components, the overall structure of the modified oligonucleotides can also be expressed in a formula to indicate a desired construction.

Thus, in one group of embodiments, the modified oligonucleotides have the formula:



wherein R^2 and R^3 represent the termini of the modified oligonucleotide; the subscript n is an integer of from 4 to 70, more preferably 4 to 25 and still more preferably 4 to 10; each B is a member independently selected from adenine, thymine, cytosine, guanine, uracil, an unsubstituted pyrazolo[3,4-d]pyrimidine and a 3-substituted pyrazolo[3,4-d]pyrimidine; and each M is selected from an oligomer-forming sugar and a peptide-nucleic acid-forming amino acid, with the proviso that at least two of the B s are selected from unsubstituted pyrazolo[3,4-d]pyrimidines and a 3-substituted pyrazolo[3,4-d]pyrimidines. In certain specific embodiments, R^2 and R^3 represent the 5'- and 3'-ends of a modified oligonucleotides wherein M is an oligomer-forming sugar (e.g., 2-deoxy- β -D-ribofuranose, β -D-ribofuranose, α -D-arabinofuranosides, α -2'-deoxyribofuranosides, 2',3'-dideoxy-3'-aminoribofuranosides, and locked sugars). For those embodiments in which the modified oligonucleotides have attached minor groove binders, reporter groups, etc., each of the R^2 and R^3 groups are meant to include functional groups suitable for attachment of linking groups and the additional functional component (MGB, fluorophore, quencher, and the like). Such functional groups include, for


example, hydroxy groups, amino groups, carboxylic acid or ester groups, phosphoric, phosphonic, or phosphinic acid or ester groups, sulfonic acid and ester groups, and the like. In other embodiments, MGBs, reporter groups and the like are attached to any of the interior base/backbone groups using conventional methodology.

5

In a related aspect, the present invention provides modified oligonucleotides comprising at least one 5-substituted pyrimidine base and at least one unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidine base. The unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidine bases useful in this aspect of the invention are essentially those that have been described above (see formulae Ia and Ib). A variety of 5-substituted pyrimidine bases are useful including those described in U.S. Patent Nos. 5,645,985 and 5,484,908.

10


In preferred embodiments, the 5-substituted pyrimidine base has the formula:



IIa

15

and the unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidine base has the formula:




Ib

20

wherein each of the X^1 , X^2 and X^3 groups is independently selected from H, OH, NH_2 and a protected amino group; and each of the R^1 and R^4 groups is independently selected from (C_1-C_{12}) heteroalkyl, (C_2-C_{12}) heteroalkenyl, (C_2-C_{12}) heteroalkynyl, $-O-(C_1-C_{12})$ alkyl, $-O-(C_2-C_{12})$ alkenyl, $-O-(C_2-C_{12})$ alkynyl, $-S-(C_1-C_{12})$ alkyl, $-S-(C_2-C_{12})$ alkenyl, $-S-(C_2-C_{12})$ alkynyl, heterocyclyl (C_1-C_{12}) alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl, aryl (C_2-C_{12}) alkenyl, aryl (C_2-C_{12}) alkynyl, aryl, heterocyclyl, halogen, $-CN$, $-CONH_2$ and protected forms thereof, with the additional feature that R^1 can also be H.

25

In particularly preferred embodiments, the 5-substituted pyrimidine base is selected from:




wherein R^4 is preferably selected from propynyl, hydroxypropynyl, aminopropynyl, or a substituted or unsubstituted phenyl, thienyl, thiazolyl, imidazolyl, furanyl, oxazolyl, pyridinyl, pyrrolyl, indolyl, benzimidazolyl, benzthiazolyl or benzoxazolyl group attached directly to the pyrimidine ring or attached to the ring via a one to four carbon linking group which can be saturated (e.g., ethylene, propylene, butylene) or unsaturated (e.g., acetylenyl, propynylene, butynylene, propenylene, butenylene).

In other preferred embodiments, the pyrazolo[3,4-d]pyrimidine base is selected from the preferred bases provided above (Ic, Id and Ie). Additionally, suitable bases are also depicted in Figure 1.

As with the earlier aspect of the invention wherein the modified oligonucleotides comprise at least two unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines, this aspect of the invention can similarly comprise additional groups such as MGBs and reporter groups (e.g., fluorophores, quenchers and the like) as well as linking groups suitable for the attachment of these additional components.

Also preferred are those embodiments in which the modified oligonucleotide is represented by the formula:




wherein R^2 represents a first end of the modified oligonucleotide; R^3 represents a second end of the modified oligonucleotide; the subscript n is an integer of from 4 to 70; each B is independently selected from adenine, thymine, cytosine, guanine, uracil, a 5-substituted pyrimidine, an unsubstituted pyrazolo[3,4-d]pyrimidine and a 3-substituted pyrazolo[3,4-d]pyrimidine (with the proviso that at least one base is a 5-substituted pyrimidine and at least one base is a pyrazolo[3,4-d]pyrimidine); and each M is an oligomer-forming sugar or a peptide-nucleic acid-forming amino acid.

In yet another aspect of the invention, modified oligonucleotides are provided which comprise an attached minor groove binder and in which as few as one of the bases is

replaced with a modified base. Surprisingly, the combination of MGBs and a single modified base (or optionally multiple modified bases) leads to modified oligonucleotides having particularly useful properties for mismatch discrimination as well as primer extensions and other utilities described below. This group of modified oligonucleotides preferably comprise from 4 to 70 bases, and an attached minor groove binder, wherein at least one of the bases is replaced by a modified base selected from the group consisting of 5-substituted pyrimidines and unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines.

In this aspect of the invention, the minor groove binders, 5-substituted pyrimidines and unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines can be essentially any of those components described above.


In one group of embodiments, the modified oligonucleotide has at least one 5-substituted pyrimidine, preferably having the formula:




IIa

wherein X^3 is selected from H, NH_2 , OH and SH; and R^4 is selected from (C_1-C_{12}) heteroalkyl, (C_2-C_{12}) heteroalkenyl, (C_2-C_{12}) heteroalkynyl, $-O-(C_1-C_{12})$ alkyl, $-O-(C_2-C_{12})$ alkenyl, $-O-(C_2-C_{12})$ alkynyl, $-S-(C_1-C_{12})$ alkyl, $-S-(C_2-C_{12})$ alkenyl, $-S-(C_2-C_{12})$ alkynyl, heterocyclyl (C_1-C_{12}) alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl, aryl (C_2-C_{12}) alkenyl, aryl (C_2-C_{12}) alkynyl, aryl, heterocyclyl, halogen, $-CN$, $-CONH_2$ and protected forms thereof. In this group of embodiments, preferred aryl and heterocyclyl group (including those components of other groups, e.g., arylalkyl) are selected from substituted and unsubstituted versions of phenyl, tolyl, pyridyl, thiazolyl, imidazolyl, furanyl, oxazolyl, thienyl, pyrrolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indolyl, triazinyl, pyrimidinyl and naphthyl. Suitable substituents for these aryl and heterocyclyl groups are those provided in the general definitions above.

More preferably, the modified oligonucleotides comprise at least one 5-substituted pyrimidine based group of Formula IId and IIe, where the R^4 substituent group is selected from a substituted or unsubstituted alkyl group or alkynyl group. Preferably, R^4 is a hydroxypropynyl, aminopropynyl or propynyl group. Alternatively, the R^4 substituent can be a 3-(hydroxymethyl)-4-hydroxy-1-butynyl group (described in more detail below).




IIId



IIe

In another group of preferred embodiments, the modified oligonucleotide has
 5 at least one modified base having the formula:




Ib

wherein each of the X^1 and X^2 groups is independently selected from H, OH, NH_2 and a
 protected amino group; and R^1 is selected from H, (C_1-C_{12}) heteroalkyl, $(C_2-$
 10 $C_{12})$ heteroalkenyl, (C_2-C_{12}) heteroalkynyl, $-O-(C_1-C_{12})$ alkyl, $-O-(C_2-C_{12})$ alkenyl, $-O-(C_2-$
 $C_{12})$ alkynyl, $-S-(C_1-C_{12})$ alkyl, $-S-(C_2-C_{12})$ alkenyl, $-S-(C_2-C_{12})$ alkynyl, heterocyclyl $(C_1-$
 $C_{12})$ alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl,
 aryl (C_2-C_{12}) alkenyl, aryl (C_2-C_{12}) alkynyl, aryl, heterocyclyl, halogen, $-CN$, $-CONH_2$ and
 protected forms thereof. In this group of embodiments, preferred aryl and heterocyclyl group
 15 (including those components of other groups, e.g., arylalkyl, heterocyclylalkyl) are selected
 from substituted and unsubstituted versions of phenyl, tolyl, pyridyl, thiazolyl, imidazolyl,
 furanyl, oxazolyl, thienyl, pyrrolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indolyl,
 triazinyl, pyrimidinyl and naphthyl. Suitable substituents for these aryl and heterocyclyl
 groups are those provided in the general definitions above.

20

In particularly preferred embodiments, the modified oligonucleotides comprise
 at least one monomer of formula If,



If


wherein X^1 and X^2 are independently H, OH or NH_2 ; and R^1 is most preferably 3-hydroxypropyn-1-yl, propynyl, 3-aminopropyn-1-yl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, halogen or 3-trifluoro-propyn-1-yl. Syntheses of some of these monomers have been reported (Balow et al., *Nuc. Acid Res.*, **26**:3350-3357 (1998); Seela et al., *J. Chem. Soc. Perkin Trans., I*, 479-488 (1999); Ramzaeva et al., *Helv. Chim. Acta*, **80**:1809-1822 (1997)).

Preparation of Modified Bases and Oligonucleotides

Reaction Schemes 1-6 provide illustrative methods for preparing a number of modified bases (unsubstituted and 3-substituted pyrazolo[3,4-d]pyrimidines and 5-substituted pyrimidines) that are useful in the present invention. The schemes illustrate the preparation of phosphoramidite derivatives of the modified bases which can be used in, for example, automatic synthesizers for preparing the modified oligonucleotides of the invention.


Reaction Scheme 1 illustrates the preparation of 5-(prop-2-ynyl-4-methylbenzoate)-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxyuridine 3'-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] (**3**) starting from 5-iodo-2'-deoxyuridine. The starting materials was treated with prop-2-ynyl 4-methylbenzoate in the presence of $Pd(PPh_3)_4$ -CuI to yield the methylbenzoate derivative (**1**) which was then converted to the 5'-blocked DMTr derivative (**2**) and then to the phosphoramidite (**3**).

Reaction Scheme 1



- Reaction Schemes 2 and 3 provide one method for the preparation of pyrazolo[3,4-d]pyrimidine phosphoramidites. More particularly, these schemes illustrate the preparation of 3-[-4((1E)-1-aza-2-methylprop-1-enyl)-1-((2R,5R)-4-{[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}-5-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}oxolan-2-yl)pyrazolo[3,4-d]pyrimidin-3-yl]prop-2-ynyl 4-methylbenzoate (**13**; $\text{R}_1 = -\text{OCOPhCH}_3$) is synthesized in two portions.

Reaction Scheme 2




In the first portion (Reaction Scheme 2), (2R,5R)-5-(4-amino-3-iodopyrazolo[3,4-d]pyrimidinyl)-2-(hydroxymethyl)oxolan-3-ol (**9**) was synthesized starting from 1,5-hydropyrazolo[3,4-d]pyrimidin-4-one. The starting material was treated with iodine monochloride and P_2S_5 successively to yield the iodo (**4**) and thione (**5**) derivatives respectively. The thione (**5**) was converted to the ethylthio-derivative (**6**), which was combined with a 1-chloro-1,2-dideoxy-3,5-di-*O*-toluoylribofuranose derivative to yield the blocked nucleoside (**7**). Reaction of compound (**7**) with sodium methoxide followed by ammonium hydroxide gave hydroxymethyl derivative (**8**) and compound (**9**) respectively.

In the second portion (Reaction Scheme 3), 2, 3-[-4((1E)-1-aza-2-methylprop-1-enyl)-1-((2R,5R)-4- {[bis(methylethyl)amino](2-cyanoethoxy)phosphinooxy}-5- {[bis(4-methoxyphenyl)phenylmethoxy]methyl} oxolan-2-yl)pyrazolo[3,4-d]pyrimidin-3-yl]prop-2-ynyl 4-methylbenzoate (**13** R_1 =-OCOPhCH₃) was prepared starting from compound (**9**).


Compound (**9**) was reacted with prop-2-ynyl 4-methylbenzoate in the presence of $Pd(PPh_3)_4$ -CuI to yield the prop-2-ynyl derivative (**10**; R_1 =-OCOPhCH₃). The amino group in this compound was protected by reaction with *N,N* dimethylacetamide dimethyl acetal to give (**11**; R_1 =-OCOPhCH₃). Compound (**11**; R_1 =-OCOPhCH₃) was converted to the DMTr derivative (**12**; R_1 =-OCOPhCH₃) and then to the phosphoramidite (**13**; R_1 =-OCOPhCH₃).

Reaction Scheme 3



- 5 Reaction Scheme 4 illustrates the preparation of N-{3-[1-((2R,5R)-5-{bis(4-methoxyphenyl)phenylmethoxy)methyl}-4-{[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}oxolan-2-yl)-6-amino-4-oxo(5-hydropyrazolo[3,4-d]pyrimidin-3-yl)]propyl}-2,2,2-trifluoroacetamide (**22**).


Reaction Scheme 4




- In Reaction Scheme 4, 6-amino-4-methoxypyrazolo[3,4-d]pyrimidine was converted to the iodo derivative (14) which was treated with 1-chloro-1,2-dideoxy-3,5-di-*O*-toluoylribofuranose to yield the nucleoside (15). Compound (15) was treated with NaOMe/MeOH to hydrolyze the toluoyl groups and yield the unblocked nucleoside (16). Treatment of (16) with aqueous sodium hydroxide gave (17) which could be converted to the

trifluoro-N-prop-2-ynyl acetamide derivative (18). Compound (18) was reduced with hydrogen and a Pd catalyst to yield the trifluoro-N-propylacetamide derivative (19). The 4-amino group of compound (19) was protected by treatment with N,N-dimethylformamide dimethylacetal to give compound (20) which was converted to DMTr derivative (21) and then to the phosphoramidite derivative (22).

Compounds of Formula 4 (3-(((2R,5R)-5-(6-amino-4-oxo-3-prop-1-ynyl(5-hydropyrazolo[3,4-d]pyrimidinyl))-2-[[bis(4-methoxyphenyl)phenylmethoxy]-methyl}oxolan-3-yloxy)[bis(methylethyl)amino]phosphinoxy]propanenitrile) (PPPG phosphoramidite) can be synthesized using some of the reactions in Reaction Scheme 4.



Formula 4




Formula 5

In the conversion of (17) to (18), prop-1-yne is used in place of 2,2,2-trifluoro-N-prop-2-ynylacetamide. The 6-amino group can be protected as described for compound (20) and the 5'-hydroxyl can be protected with a DMTr group as described for compound (21). Finally the phosphoramidite can be prepared as described for compound (22) to yield the compound of Formula 4. In a similar way compounds of Formula 5 where R⁶ is -OCOPhCH₃ or -NHCOCF₃ can be prepared using related reactions known in the art.

Reaction Scheme 5 illustrates the preparation of 3-{[5-(4,6-bis{(1E)-1-aza-2-[[bis(2-methylpropyl)amino]vinyl]-3-prop-1-ynylpyrazolo[3,4-d]pyrimidinyl))-2-[[bis(4-methoxyphenyl)phenylmethoxy]-methyl}oxolan-3-yloxy][ethyl(methylethyl)amino]-phosphino}propanenitrile (26).


Reaction Scheme 5



Compound **(26)** can be synthesized starting from $(\text{NH}_2)_2\text{PPPA}$ (**(23)**) (see, Seela & Driller, *Helv. Chim. Acta* **71**:757-761(1988)). Compound **(23)** can be converted to the bis(methylethyl)amino derivative **(24)** (Vincent et al, *J. Org. Chem.*, **64**:991-997 (1999)), followed by reaction first with DMTrCl to yield **(25)** that can be converted to the phosphoramidite **(26)**.


Reaction Scheme 6 provides the synthesis of protected 3-substituted pyrazolo[3,4-d]pyrimidines in which the substituent is a heteroalkyl group.

Reaction Scheme 6




- Reaction Scheme 7 provides the synthesis of protected 3-substituted pyrazolo[3,4-d]pyrimidine in which the substituent is a methoxypropynyl group. The 3-iodo diamino derivative (**23**) was reacted with $\text{Pd(PPh}_3)_4\text{-CuI}$, triethylamine in anhydrous DMF followed by methyl propargyl ether to yield the 3-methoxypropynyl-2,4-diaminopyrazolo[3,4-d]pyrimidine derivative (**27**). The amino groups in this compound were blocked by reaction with N,N-dimethylformamide dimethylacetal to yield (**28**). The blocked nucleoside was first reacted with dimethoxytrityl chloride and then with 2-cyanoethyl diisopropylchlorophosphoramidite to yield the desired blocked phosphoramidite (**29**).

Reaction Scheme 7



The following reaction schemes provide methods for the preparation of 3-heterocycle substituted pyrazolo[3,4-d]pyrimidines. The general methods provided herein can be adapted for the preparation of other heterocyclic substituents.


Reaction Scheme 8



Reaction Scheme 8 provides the synthesis of protected 3-substituted pyrazolo[3,4-d]pyrimidine in which the substituent is a furanyl group. Malonitrile was reacted with 2-furfuryl chloride in the presence of a base, followed by treatment with dimethylsulfide to give the methoxy dinitrile derivative (30). Reaction of (30) with hydrazine yielded the substituted pyrazole (31) which was reacted with formamide to give 3-(2-furyl)pyrazolo[3,4-d]pyrimidine-4-ylamine (32). Base (32) was treated with 1-chloro-1,2-dideoxy-3,5-di-O-toluoylribofuranose to yield the blocked nucleoside (33). Compound (33) was first treated with NaOMe/MeOH to hydrolyze the toluoyl groups to yield the unblocked nucleoside which was reacted with N,N-dimethylformamide dimethylacetal to yield the protected nucleoside derivative (34). This derivative was first reacted with dimethoxytrityl

chloride and then with 2-cyanoethyl diisopropylchlorophosphoramidite to yield the desired blocked phosphoramidite (**35**).

Reaction Scheme 9



5

Reaction Scheme 9 provides the synthesis of protected 3-substituted 2,4-diamino pyrazolo[3,4-d]pyrimidine in which the substituent is a furanyl group. The carbonitrile (**31**) was reacted with guanidinium carbonate to yield 3-(2-furyl)pyrazolo[3,4-d]pyrimidine-4,6-diamine (**36**). This base was treated with 1-chloro-1,2-dideoxy-3,5-di-O-toluoylribofuranose to yield the blocked nucleoside which was treated with NaOMe/MeOH to hydrolyze the toluoyl groups and yield the unblocked nucleoside derivative (**37**). The latter compound was reacted with N,N-dimethylformamide dimethylacetal to yield the protected nucleoside derivative (**38**). This derivative was first reacted with dimethoxytrityl chloride and then with 2-cyanoethyl diisopropylchlorophosphoramidite to yield the desired blocked phosphoramidite (**39**).

15


In each of the above schemes, one of skill in the art will understand that other protecting groups and/or activating groups can be used. Additionally, different non-natural

base-containing monomers, with different functional groups, can require different blocking groups for successful synthesis of the modified oligonucleotides. A variety of useful protection groups, their synthesis and de-protection methods are described in, for example, Beaucage & Iyer, *Tetrahedron* **48**:2223-2311 (1992).

5 The synthesis of oligonucleotides, and modified oligonucleotides can be initiated from a solid support containing a cleavable linker to which is attached the first base. The oligonucleotides of the invention can be synthesized containing a substituted pyrazolo[3,4-d]pyrimidine as the first nucleotide at the 3'-end, using an appropriate 3-substituted pyrazolo[3,4-d]pyrimidine. Alternatively, a 5-substituted pyrimidine can be
10 attached initially to a solid support. Reaction Scheme 10 illustrates a general approach to the synthesis of a protected 3-substituted pyrazolo[3,4-d]pyrimidine dA CPG derivative (C). Intermediate (A) can be prepared using procedures described in Reaction Scheme 3, while (B) and (C) can be synthesized as described in co-pending U.S. application Ser. No. 09/457,616.

15

Reaction Scheme 10



In this aspect of the invention, a variety of solid supports are useful, provided the support is compatible with automated oligonucleotide techniques and include, glass, polystyrene, nylon, plastic and the like. Additionally, the present invention provides in a related aspect, a solid support (e.g., controlled pore glass (CPG)) having an attached modified base, preferably those of formulae Ic, Id, Ie, If, IIb, IIc, IId, IIe. More preferably, the attached modified base is a 3-substituted pyrazolo[3,4-d]pyrimidine or a 5-substituted pyrimidine moiety wherein the substituents are selected from the preferred groups described above, and further including the novel bases described in more detail below.

The present invention further includes those compositions and methods wherein any of the unsubstituted and/or 3-substituted pyrazolo[3,4-d]pyrimidines and/or 5-substituted pyrimidines are combined or used in combination with other modified bases known in the art. Other modified monomeric units have been disclosed Scheit,

NUCLEOTIDE ANALOGS, John Wiley, New York, 1980; Uhlman and Meyman, *Chemical Reviews*, **90**:543-584 (1990), Seela and Debelak, *Nucl. Acids Res.*, **28**:3224-3232 (2000); Balow et al, *Nucl. Acids Res.*, **26**:3350-3357 (1998); Bolli et al *Nucl. Acids Res.*, **24**:4660-4667 (1996).

5

Modified Oligonucleotide Arrays

In another embodiment of the present invention, modified oligonucleotides are used in procedures which utilize arrays of oligonucleotides, such as sequencing by hybridization and array-based analysis of gene expression. A variety of arrays are contemplated by the present invention including, for example, chip or platform arrays, bead arrays, liquid phase arrays, 'zip-code' arrays and the like. In sequencing by hybridization, an ordered array of oligonucleotides of different known sequences is used as a platform for hybridization to one or more test polynucleotides, nucleic acids or nucleic acid populations. Determination of the oligonucleotides which are hybridized and alignment of their known sequences allows reconstruction of the sequence of the test polynucleotide. Alternatively, oligonucleotides comprising the wild-type sequence and all possible mutant sequences for a given region of a gene of interest can be placed on an array. Exposure of the array to DNA or RNA from a subject or biological specimen, under hybridization conditions, allows determination of wild-type or mutant status for the gene of interest. See, for example, U.S. Patent Nos. 5,492,806; 5,525,464; 5,556,752; and PCT Publications WO 92/10588 and WO 96/17957. Both of these techniques require discrimination between related sequences, especially at the single-nucleotide level; hence, the enhanced discriminatory properties of the modified oligonucleotides of the invention will provide improvements in these techniques. Materials for construction of arrays include, but are not limited to, nitrocellulose, glass, silicon wafers, optical fibers and other materials suitable for construction of arrays such as are known to those of skill in the art. The synthesis of oligonucleotides arrays has been described in co-pending U.S. application Ser. No. 09/364,320, and suitable modifications can be made to the methods therein for preparing the arrays of the present invention.

An additional application of the present invention to array technology is in the examination of patterns of gene expression in a particular cell or tissue. In this case, oligonucleotides or polynucleotides corresponding to different genes are arrayed on a surface, and a nucleic acid sample from a particular cell or tissue type, for example, is incubated with the array under hybridization conditions. Detection of the sites on the array at which

hybridization occurs allows one to determine which oligonucleotides have hybridized, and hence which genes are active in the particular cell or tissue from which the sample was derived.

Array methods can also be used for identification of mutations or

5 polymorphisms, where wild-type and mutant sequences are placed in an ordered array on a surface. Hybridization of a polynucleotide sample to the array under stringent conditions, and determination of which oligonucleotides in the array hybridize to the polynucleotide, allows determination of whether the polynucleotide possesses the wild-type or the mutant sequence. Since many mutant sequences of clinically-relevant genes differ from their wild-
10 type counterpart at only one or a few nucleotide positions, the enhanced discriminatory powers of the modified oligonucleotides of the invention will provide improvements in mutation detection.

In all of the above-mentioned applications of array technology, the increased discriminatory abilities of modified oligonucleotide provide significant improvements in
15 sensitivity and resolving power.

Efficient hybridization, for example in arrays, requires that the capture probes contain T_m 's in a narrow temperature range. In one embodiment of the invention the T_m of capture oligonucleotides for use in arrays is balanced, eliminating the problem associated with GC and AT-rich sequences, by using a combination of 3'-MGBs and modified bases
20 (e.g., PPPA and PPPU). Table 1 and Figure 2 illustrate the balancing of the T_m of pyrimidine rich probes with different G/C and A/T base pairs. To balance the T_m of the different probes, PPPA, PU and MGB were used in combination. The purine and pyrimidine-rich probes and complementary targets are shown in Table 1 and the T_m s of these probes are shown in Figure 2. As shown in Figure 2 the T_m difference between a GC-rich and an AT-
25 rich probe can be more than 50°C.

Table 1

Sequences of Oligonucleotide Probes and Complementary Targets

	<u>Complementary Targets</u>	<u>Py/Pu-rich probes</u>	<u>base pairs</u>
1*	TCGGCGGCGT	1*.MGB-Q-CGCCGCCG	8 G/C
2*	ACAGCGGCGT	2*.MGB-Q-CGCCGCTG	7 G/C, 1 A/T
3*	ACAGCGACGT	3*.MGB-Q-CGTCGCTG	6 G/C, 2 A/T
4*	TCAGTGACGA	4*.MGB-Q-CGTCACTG	5 G/C, 3 A/T
5*	TCAGTGACAA	5*.MGB-Q-TGTCACTG	4 G/C, 4 A/T

6*	TCAATGACAG	6*.MGB-Q-TGTCATTG	3 G/C, 5 A/T
7*	ACAATGATAA	7*.MGB-Q-TATCATTG	2 G/C, 6 A/T
8*	CCAATAATAA	8*.MGB-Q-TATTATTG	1 G/C, 7 A/T
9*	GTAATAATAA	9*.MGB-Q-TATTATTA	8 A/T


As can be seen in Figure 2, modification with PPPA, PU and a MGB provides a probe set having balanced T_m s.

Efficient hybridization for example in arrays, requires that the probes contain T_m 's in a narrow temperature range. Thus, in related embodiments, the modified oligonucleotides described herein can comprise any combination of normal bases, unsubstituted pyrazolo[3,4-d]pyrimidine bases (e.g., PPG and PPA), 3-substituted pyrazolo[3,4-d]pyrimidines, modified purine, modified pyrimidine, 5-substituted pyrimidines, universal bases and a minor groove binder to balance the T_m of the oligonucleotide.

The ability to predict and "level" the T_m s of multiple modified oligonucleotides is particularly useful for the development of oligonucleotide arrays or for compositions having a plurality of oligonucleotides. Still further, the ability to predict T_m s for modified oligonucleotides is useful for establishing appropriate conditions for hybridization, renaturation, mapping variations of base compositions of sequences, determination of sequence complexity and divergence. A T_m prediction algorithm can be developed using models that assign thermodynamic contributions to duplex formation for all possible individual base pair nearest neighbors. Thermodynamic parameters for 10 independent nearest-neighbors of base pairs are available from published sources for DNA:RNA hybrids (see Sugimoto et al., *Biochem.* **34**:11211-11216 (1995)) and dsDNA (SantaLucia et al, *Biochem.* **36**:10581-10595 (1997)). The use of the nearest neighbor parameters to develop programs to calculate oligonucleotide duplex stability in terms of T_m have been described (see, Schutz & von Ahsen, *Biotechniques*, **27**:1218-22 (1999); Owczarzy et al, *Biopolymers*, **44**:217-239 (1997); for PNA/DNA, see Geisen et al, *Nucl. Acids Res.*, **26**:5004-5006 (1998); Blake et al, *Bioinformatics*, **15**:370-5 (1999) and U.S. Patent No. 5,556,749.

The basic principles of thermodynamic and kinetic description of mismatch discrimination are well known and are summarized below.

A probe–target interaction is an equilibrium reaction that can be described as



At the temperature T equilibrium of forward (association) and reverse (dissociation) reactions can be described by equilibrium constant K where :

$$K = \frac{[AB]}{[A][B]} ;$$

in terms of equilibrium thermodynamics

$$\begin{aligned} K &= \exp(-\Delta H_{AB}^{\circ} / RT + \Delta S_{AB}^{\circ} / R); \\ \Delta G_{AB}^{\circ} &= \Delta H_{AB}^{\circ} - T\Delta S_{AB}^{\circ}; \\ \Delta G_{AB}^{\circ} &= -RT \ln(K) \end{aligned}$$

where : ΔH_{AB}° , ΔS_{AB}° and ΔG_{AB}° - enthalpy, entropy and free energy changes for duplex AB formation.

The duplex AB has N base pairs (N-1 nearest neighbors) and the nearest-neighbor approximation assumes


$$\begin{aligned} \Delta H_{AB}^{\circ} &= \Delta H_1^{\circ} + \Delta H_2^{\circ} + \dots \Delta H_{n-1}^{\circ} + \Delta H_{init}^{\circ} = \sum (\Delta H_i^{\circ} + \Delta H_{init}^{\circ}) \\ \Delta S_{AB}^{\circ} &= \Delta S_1^{\circ} + \Delta S_2^{\circ} + \dots \Delta S_{n-1}^{\circ} + \Delta S_{init}^{\circ} = \sum (\Delta S_i^{\circ} + \Delta S_{init}^{\circ}) \\ \Delta G_{AB}^{\circ} &= \Delta G_1^{\circ} + \Delta G_2^{\circ} + \dots \Delta G_{n-1}^{\circ} + \Delta G_{init}^{\circ} = \sum (\Delta G_i^{\circ} + \Delta G_{init}^{\circ}) \end{aligned}$$

where : ΔH_i° , ΔS_i° and ΔG_i° - enthalpy, entropy and free energy changes for each nearest-neighbor base pairs increment for duplex AB formation, ΔH_{init}° , ΔS_{init}° and ΔG_{init}° - thermodynamic parameters of initiation of duplex AB formation.

In the case of total concentration equality, $C_{\text{probe}} = C_{\text{target}}$, an equation for T_m in terms of free energy, enthalpy and entropy can be arrived at as shown below:

$$T_m = \frac{\sum (\Delta H_i^{\circ} + \Delta H_{init}^{\circ})}{\sum (\Delta S_i^{\circ} + \Delta S_{init}^{\circ}) + R \ln[(C_{\text{probe}} + C_{\text{target}})/4]}$$

Similarly as shown above relationships can be developed when the probe hybridizes to a target that contains a mismatch, as shown below:



Thermodynamic parameters for mismatched duplex AB^* are: K^* , $\Delta H^{\circ}_{AB^*}$, $\Delta S^{\circ}_{AB^*}$ and $\Delta G^{\circ}_{AB^*}$. In terms of nearest-neighbor parameters for a mismatched base pair at position “4”, for example, involving nearest-neighbor “3” and “4” the discrimination parameters $\Delta\Delta H^{\circ}$, $\Delta\Delta S^{\circ}$, $\Delta\Delta G^{\circ}$ can be expressed as:

$$\Delta\Delta H^{\circ} = \Delta H^{\circ}_{AB} - \Delta H^{\circ}_{AB^*} = (\Delta H^{\circ}_3 - \Delta H^{\circ*}_3) + (\Delta H^{\circ}_4 - \Delta H^{\circ*}_4) = \Delta\Delta H^{\circ}_3 + \Delta\Delta H^{\circ}_4.$$

$$\Delta\Delta S^{\circ} = \Delta S^{\circ}_{AB} - \Delta S^{\circ}_{AB^*} = (\Delta S^{\circ}_3 - \Delta S^{\circ*}_3) + (\Delta S^{\circ}_4 - \Delta S^{\circ*}_4) = \Delta\Delta S^{\circ}_3 + \Delta\Delta S^{\circ}_4.$$

$$\Delta\Delta G^{\circ} = \Delta G^{\circ}_{AB} - \Delta G^{\circ}_{AB^*} = (\Delta G^{\circ}_3 - \Delta G^{\circ*}_3) + (\Delta G^{\circ}_4 - \Delta G^{\circ*}_4) = \Delta\Delta G^{\circ}_3 + \Delta\Delta G^{\circ}_4.$$

In one embodiment of the invention, probe sequences are designed to determine the thermodynamic nearest-neighbor parameters for oligonucleotides containing normal purines, pyrimidines and a minor groove binder CDPI₃ attached to the 3'-end of an oligonucleotide. In another embodiment thermodynamic nearest-neighbor parameters for oligonucleotide containing normal pyrimidines, purines, a 3'-end attached CDPI₃, but wherein the guanine is replaced with 5-hydropyrazolo[3,4-d]pyrimin-4-one. These nearest-neighbor thermodynamic parameters are used in these two embodiments to calculate the T_m s of these types of oligonucleotides with and without CDPI₃. In the case where the CDPI₃ was attached at base 1 at the 3'-end the new T_m prediction algorithm and formula were used as derived below.

$$T_M = \frac{\sum (\Delta H^{\circ}_i + \Delta H^{\circ}_{init})}{\sum (\Delta S^{\circ}_i + \Delta S^{\circ}_{init}) + R \ln[(C_{probe} + C_{target})/4] + \Delta\Delta S^{\circ}_{MGB}}$$

The model assumes that a MGB attached to an oligonucleotide has an additional pure entropic factor $\Delta\Delta S^{\circ}_{MGB}$. This value can be calculated according to algorithm : 6 bases (5 nearest-neighbor) from 3'-prime end are covered by the MGB from starting base “1” or “2”. Each nearest-neighbors covered has unique value $\Delta\Delta S^{\circ}_{iMGB}$. Correction factor “A” are introduced in the case when adenosine can be found in base positions 6, 7 or 8. Statistical factor “n” is introduced additionally to factor “A” were n=3 for

-AAA-; n=2 for -AA-, -ANA-; n=1 for -A-; n=0 for -AN- and -NA- (N is any base other than A). A final value $\Delta\Delta S^\circ_{MGB}$ can be determined by the equation

$$\Delta\Delta S^\circ_{MGB} = \sum \Delta\Delta S^\circ_{iMGB} + A(\text{optional}) + R \cdot \ln(N) (\text{optional})$$

5

Tables 2a and 2b contain the nearest-neighbor thermodynamic parameters for oligonucleotides containing PPG alone and the entropic contribution of the MGB, respectively. In the latter case the contribution of fluorescein and a dark quencher is incorporated in the thermodynamic treatment and has generally been shown to be negligible.

10 [The quencher is part of the linker between the MGB and the oligonucleotide]

Tables 2a and 2b

- a) Nearest-neighbor parameters for PPG containing oligonucleotides
b) The entropic contribution of the CDPI₃.

15

a)

Number	5'-3'	ΔH°	ΔS°	$\Delta G^\circ(65)$
1	AA	-7850	-22.3	-324
2	AT	-8180	-23.0	-411
3	AC	-8450	-22.6	-795
4	AG	-6560	-17.2	-730
5	TA	-7230	-21.8	129
6	TT	-7850	-22.3	-324
7	TC	-8720	-24.0	-614
8	TG	-6900	-18.3	-715
9	CA	7530	-20.3	-652
10	CT	-6390	-17.3	-548
11	CC	-7860	-20.1	-1071
12	CG	-6030	-14.0	-1288
13	GA	-10070	-27.4	-795
14	GT	-9110	-23.5	-1175
15	GC	-13170	-34.0	-1665
16	GG	-8080	-20.2	-1258
17	GCinit	-48440	-17.6	1116
18	ATinit	-1060	-7.1	1341

b)

Number	5'-3'	$\Delta\Delta S^\circ$
1	AA	3.408
2	AT	3.060
3	AC	1.442
4	AG	0.750
5	TA	2.463
6	TT	3.313
7	TC	2.870
8	TG	0.893
9	CA	0.607

10	CT	2.253
11	CC	0.905
12	CG	-0.721
13	GA	1.282
14	GT	2.397
15	GC	2.172
16	GG	1.298

Use of Modified Oligonucleotides

5 The modified oligonucleotides of the present invention provides numerous advantages over unmodified oligonucleotides, including superior mismatch discrimination. The modified oligonucleotides of the invention are particularly useful as probes, wherein their hybridization to a target sequence is detected, or as primers, wherein their hybridization to a target sequence is followed by polynucleotide synthesis initiated from the 3' terminus of
10 the modified oligonucleotide, and the synthesized product (*i.e.*, the extension product) is detected.

 The modified oligonucleotides of the present invention are useful in other techniques in which hybridization of an oligonucleotide to another nucleic acid is involved. These include, but are not limited to, techniques in which hybridization of an oligonucleotide
15 to a target nucleic acid is the endpoint; techniques in which hybridization of one or more oligonucleotides to a target nucleic acid precedes one or more polymerase-mediated elongation steps which use the oligonucleotide as a primer and the target nucleic acid as a template; techniques in which hybridization of an oligonucleotide to a target nucleic acid is used to block extension of another primer; techniques in which hybridization of an
20 oligonucleotide to a target nucleic acid is followed by hydrolysis of the oligonucleotide to release an attached label; and techniques in which two or more oligonucleotides are hybridized to a target nucleic acid and interactions between the multiple oligonucleotides are measured. Conditions for hybridization of oligonucleotides, and factors which influence the degree and specificity of hybridization, such as temperature, ionic strength and solvent
25 composition, are well-known to those of skill in the art. *See*, for example, Sambrook *et al.*, *supra*; Ausubel, *et al.*, *supra*; M.A. Innis *et al.* (eds.) PCR Protocols, Academic Press, San Diego, 1990; B.D. Hames *et al.* (eds.) Nucleic Acid Hybridisation: A Practical Approach, IRL Press, Oxford, 1985; and van Ness *et al.* (1991) *Nucleic Acids Res.* **19**:5143-5151.

Hybridization of probes and/or primers to target sequences proceeds according to well-known and art-recognized base-pairing properties, such that adenine base-pairs with thymine or uracil, and guanine base-pairs with cytosine. The property of a nucleotide that allows it to base-pair with a second nucleotide is called complementarity. Thus, adenine is complementary to both thymine and uracil, and *vice versa*; similarly, guanine is complementary to cytosine and *vice versa*. An oligonucleotide which is complementary along its entire length with a target sequence is said to be perfectly complementary, perfectly matched, or fully complementary to the target sequence, and *vice versa*. An oligonucleotide and its target sequence can have related sequences, wherein the majority of bases in the two sequences are complementary, but one or more bases are noncomplementary, or mismatched. In such a case, the sequences can be said to be substantially complementary to one another. If the sequences of an oligonucleotide and a target sequence are such that they are complementary at all nucleotide positions except one, the oligonucleotide and the target sequence have a single nucleotide mismatch with respect to each other.

The modified pyrazolo[3,4-d]pyrimidine nucleotides of the invention retain the base-pairing specificity of their naturally-occurring analogues; PPPG analogues are complementary to cytosine, while PPPA analogues are complementary to thymine and uracil. The PPPG and PPPA analogues not only have a reduced tendency for so-called "wobble" pairing with non-complementary bases, compared to guanine and adenine, but the 3-substituted groups increase binding affinity in duplexes. Similarly, modified pyrimidines hybridize specifically to their naturally occurring counter partners.

Conditions for hybridization are well-known to those of skill in the art and can be varied within relatively wide limits. Hybridization stringency refers to the degree to which hybridization conditions disfavor the formation of hybrids containing mismatched nucleotides, thereby promoting the formation of perfectly matched hybrids or hybrids containing fewer mismatches; with higher stringency correlated with a lower tolerance for mismatched hybrids. Factors that affect the stringency of hybridization include, but are not limited to, temperature, pH, ionic strength, concentration of organic solvents such as formamide and dimethylsulfoxide and chaotropes. As is well known to those of skill in the art, hybridization stringency is increased by higher temperatures, lower ionic strengths, and lower solvent concentrations. See, for example, Ausubel *et al.*, *supra*; Sambrook *et al.*, *supra*; M.A. Innis *et al.* (eds.) PCR Protocols, Academic Press, San Diego, 1990; B.D. Hames *et al.* (eds.) Nucleic Acid Hybridisation: A Practical Approach, IRL Press, Oxford, 1985; and van Ness *et al.*, (1991) *Nucleic Acids Res.* **19**:5143-5151.

Thus, in the formation of hybrids (duplexes) between an oligonucleotide and its target sequence, the oligonucleotide is incubated in solution, together with a polynucleotide containing the target sequence, under conditions of temperature, ionic strength, pH, *etc.*, that are favorable to hybridization, *i.e.*, under hybridization conditions.

5 Hybridization conditions are chosen, in some circumstances, to favor hybridization between two nucleic acids having perfectly-matched sequences, as compared to a pair of nucleic acids having one or more mismatches in the hybridizing sequence. In other circumstances, hybridization conditions are chosen to allow hybridization between mismatched sequences, favoring hybridization between nucleic acids having fewer mismatches.

10 The degree of hybridization of an oligonucleotide to a target sequence, also known as hybridization strength, is determined by methods that are well-known in the art. A preferred method is to determine the T_m of the hybrid duplex. This is accomplished, as described *supra*, by subjecting a duplex in solution to gradually increasing temperature and monitoring the denaturation of the duplex, for example, by absorbance of ultraviolet light,
15 which increases with the unstacking of base pairs that accompanies denaturation. T_m is generally defined as the temperature midpoint of the transition in ultraviolet absorbance that accompanies denaturation. Alternatively, if T_m s are known, a hybridization temperature (at fixed ionic strength, pH and solvent concentration) can be chosen that it is below the T_m of the desired duplex and above the T_m of an undesired duplex. In this case, determination of
20 the degree of hybridization is accomplished simply by testing for the presence of hybridized probe.

If a probe comprises a detectable label, assays for hybridized probe are usually designed to detect the presence of label in duplex material. This can be accomplished, for example, by specifically selecting duplex material, specifically destroying single-stranded
25 material, or utilizing some combination of these methods. For example, hybridization reaction mixtures can be subjected to high-stringency conditions and/or single strand-specific nucleases; or duplexes can be purified by affinity techniques specific for double-stranded, as opposed to single-stranded, nucleic acids. In a preferred embodiment of the invention, duplexes are detected by release of label from a probe under conditions in which label is
30 released only when the probe is in a duplex. Another embodiment requires the separation of the label and quenchers when hybridized to the target.

Detectable labels or tags suitable for use with nucleic acid probes are well-known to those of skill in the art and include, but are not limited to, radioactive isotopes, chromophores, fluorophores, chemiluminescent and electrochemiluminescent agents,

magnetic labels, microspheres, colloidal metal (Taton et al, *Science* **289**:1757-1760 (2000)), immunologic labels, ligands and enzymatic labels. Suitable labels also include mass labels and those used in deconvolution of combinatorial chemistry libraries, for example, tags that can be recognized by high performance liquid chromatography (HPLC), gas chromatography, mass spectrometry, optical imaging fibers, surface plasmon resonance, correlation spectroscopy, nanotechnology (Guetence et al, *J. Chromatogr. B. Biomed. Sci. Appl.* **739**:139-150 (2000)) and the like.

Methods for labeling of oligonucleotides are well-known to those of skill in the art and include, for example, chemical and enzymatic methods. By way of example, methods for incorporation of reactive chemical groups into oligonucleotides, at specific sites, are well-known to those of skill in the art. Oligonucleotides containing a reactive chemical group, located at a specific site, can be combined with a label attached to a complementary reactive group (*e.g.*, an oligonucleotide containing a nucleophilic reactive group can be reacted with a label attached to an electrophilic reactive group) to couple a label to a probe by chemical techniques. Exemplary labels and methods for attachment of a label to an oligonucleotide are described, for example, in U.S. Patent No. 5,824,796; U.S. Patent No. 5,210,015; Kessler (ed.), *Nonradioactive Labeling and Detection of Biomolecules*, Springer-Verlag, Berlin, 1992; Kricka (ed.) *Nonisotopic DNA Probe Techniques*, Academic Press, San Diego, 1992; Howard (ed.) *Methods in Nonradioactive Detection*, Appleton & Lange, Norwalk, 1993. Non-specific chemical labeling of an oligonucleotide can be achieved by combining the oligonucleotide with a chemical that reacts, for example, with a particular functional group of a nucleotide base, and simultaneously or subsequently reacting the oligonucleotide with a label. See, for example, Draper *et al.* (1980) *Biochemistry* **19**:1774-1781. Enzymatic incorporation of label into an oligonucleotide can be achieved by conducting enzymatic modification or polymerization of an oligonucleotide using labeled precursors, or by enzymatically adding label to an already-existing oligonucleotide. See, for example, U.S. Patent No. 5,449,767. Examples of modifying enzymes include, but are not limited to, DNA polymerases, reverse transcriptases, RNA polymerases, *etc.* Examples of enzymes which are able to add label to an already-existing oligonucleotide include, but are not limited to, kinases, terminal transferases, ligases, glycosylases, *etc.*

If an oligonucleotide is capable of acting as a primer, the degree of hybridization of the oligonucleotide can also be determined by measuring the levels of the extension product of the primer. In the case, either the primer can be labeled, or one or more of the precursors for polymerization (normally nucleoside triphosphates) can be labeled.

Extension product can be detected, for example, by size (e.g., gel electrophoresis), affinity methods, or any other technique known to those of skill in the art.

Primer extension (“minisequencing”, “genetic bit analysis”) assays are commonly used for SNP typing and have the potential to be used in other genotyping and mutation screening applications (Pastinen T. et al., *Genome Res.*, **10**:1031-42 (2000)). In certain embodiments modified bases and minor groove binders improve primer extension assays in several ways. The added duplex stability provided by MGB, or 5-substituted pyrimidine or 3-substituted pyrazolo[3,4-d]pyrimidine enables extensions to be performed at elevated temperatures. This is advantageous because problematic secondary structures in target molecules are eliminated at elevated temperatures. Also, hybridization of target to primer is faster at higher temperature. Thermostable polymerases such as Taq polymerase and Bst DNA polymerase may be used in such reactions.

Furthermore, MGBs and modified bases improve the specificity of assays by eliminating one class of false positive signals. Primer sequences that form hairpin structures or homodimers are prone to template-independent extension (the 5’ end of the primer functions as template), resulting in false positive signal. MGBs and modified bases on “templates” inhibit extension by DNA polymerases. Thus, MGBs on the 5’ end, or modified bases on the 5’ end or middle of a primer, can prevent extension (false positives) from primer hairpins or primer dimers. Finally, PPG can be used to eliminate non-canonical structures formed by G-rich oligonucleotides, enabling primer extension assays in such sequences.

Other assays in which the present modified oligonucleotides are particularly useful are described in co-pending application Ser. No. 09/054,832.

Still other amplification assays in which modified oligonucleotides are useful include the amplification assays based on the invasive cleavage of oligonucleotide probes by flap endonucleases (Lyamichev et al., *Nature Biotechnol.*, **17**:292-296 (1999)); self-sustained sequence replication type assays (Mueller et al, *Histochem. Cell Biol.*, **108**:431-437 (1997)) and the like. Surprisingly, non-natural bases can be substituted in both the invader and genomic probes of a cleavase-based assay. These modifications include but are not limited to pyrazolo[3,4-d]pyrimidines, 3-substituted pyrazolo[3,4-d]pyrimidines and 5-substituted pyrimidines. Non-natural backbones are also included such as monomers used in peptide nucleic acids, locked nucleic acids etc. Still further, the modified oligonucleotides can have attached minor groove binders, fluorophores, quenchers, and the like. One of skill in the art will appreciate that chimeras can also be used to allow optimal enzyme activity and performance.

In the present invention, marked improvements are unexpectedly shown in cleavase-based assays, when modified bases were substituted for normal bases in both the invader and genomic probes. Thus, the use of the modified bases hydroxypropynylPPA (HOPPPA) and 3-iododiaminoPPA allowed one to decrease the length of both the invader and genomic probes, yet provide improved assay performance. In another embodiment non-natural bases are also incorporated in the cassette probe.

In view of the above, the present invention provides in one aspect, a method for distinguishing polynucleotides with related sequences, the method comprising:

(a) contacting a modified oligonucleotide having a defined sequence comprising at least one 3-substituted pyrazolo[3,4-*d*]pyrimidine or 5-substituted pyrimidine in place of a purine or pyrimidine base with at least two polynucleotides, wherein one of the polynucleotides has a target sequence that is perfectly complementary to the modified oligonucleotide and at least one of the other polynucleotides has a target sequence with at least one base mismatch; and

(b) determining the degree of hybridization between the modified oligonucleotide and each of the polynucleotides.

Preferably, at least one of the other polynucleotides has a target sequence with one or two base mismatches, more preferably only one base mismatch.

As noted above, a target sequence refers to a nucleotide sequence which comprises a site of hybridization for a probe or a primer. Target sequences can be found in any nucleic acid including, but not limited to, genomic DNA, cDNA, RNA and any amplified product thereof, and can comprise a wild-type gene sequence, a mutant gene sequence, a non-coding sequence, a regulatory sequence, *etc.* A target sequence will generally be less than 100 nucleotides, preferably less than 50 nucleotides, and most preferably, less than 21 nucleotides in length.

The modified oligonucleotides used in this aspect of the invention are essentially modified probes and the polynucleotides can be distinguished by determining which polynucleotides hybridizes to the modified probe. The modified probes can be labeled with any detectable label, or the probe can have the capacity to become labeled either before or after hybridization, such as by containing a reactive group capable of association with a label or by being capable of hybridizing to a secondary labeled probe, either before or after hybridization to the target. Conditions for hybridization of nucleic acid probes are well-known to those of skill in the art. *See, for example, Sambrook et al., supra; Ausubel et al., supra; Innis et al., supra; Hames et al., supra; and van Ness et al., supra.*

Hybridization can be assayed (*i.e.*, hybridized nucleic acids can be identified) by distinguishing hybridized probe from free probe by one of several methods that are well-known to those of skill in the art. These include, but are not limited to, attachment of target nucleic acid to a solid support, either directly or indirectly (by hybridization to a second, support-bound probe or interaction between surface-bound and probe-conjugated ligands) followed by direct or indirect hybridization with probe, and washing to remove unhybridized probe; determination of nuclease resistance; buoyant density determination; affinity methods specific for nucleic acid duplexes (*e.g.*, hydroxyapatite chromatography); interactions between multiple probes hybridized to the same target nucleic acid; *etc.* See, for example, Falkow *et al.*, U.S. Patent No. 4,358,535; Urdea *et al.*, U.S. Patent Nos. 4,868,105 and 5,124,246; Freifelder, *Physical Biochemistry*, Second Edition, W. H. Freeman & Co., San Francisco, 1982; Sambrook, *et al.*, *supra*; Ausubel *et al.*, *supra*; Hames *et al.*, *supra*; and other related references. The duplex-stabilizing capability of MGB-modified oligonucleotide conjugates makes hybridization possible under more stringent conditions, wherein potentially occluding secondary structure in the target nucleic acid can be minimized. Accordingly, such MGB-modified oligonucleotides are particularly preferred in this aspect of the invention.

In a related aspect, the present invention provides a method for detecting the presence of a target sequence in a polynucleotide, the method comprising:

(a) incubating a polynucleotide to be tested for the presence of the target sequence with a modified oligonucleotide having a sequence that is substantially complementary to the target sequence under hybridization conditions; and

(b) identifying hybridized nucleic acids;

wherein the modified oligonucleotide comprises at least one 3-substituted pyrazolo[3,4-*d*]pyrimidine in place of a purine residue.

Preferably, the modified oligonucleotide is a labeled probe and has at least two 3-substituted pyrazolo[3,4-*d*]pyrimidine bases. In this group of embodiments, a labeled probe is hybridized to a target and/or an extension product of a target, and a change in the physical state of the label is effected as a consequence of hybridization. As used herein, a "probe" is a nucleic acid molecule that is capable of hybridizing to a target sequence in a second nucleic acid molecule. By way of example, one assay of this type, the hydrolyzable probe assay, takes advantage of the fact that many polymerizing enzymes, such as DNA polymerases, possess intrinsic 5'-3' exonucleolytic activities. Accordingly, if a probe is hybridized to a sequence that can serve as a template for polymerization (for instance, if a probe is hybridized to a region of DNA located between two amplification primers, during

the course of an amplification reaction), a polymerizing enzyme that has initiated polymerization at an upstream amplification primer is capable of exonucleolytically digesting the probe. Any label attached to such a probe will be released, if the probe is hybridized to its target and if amplification is occurring across the region to which the probe is hybridized.

5 Released label is separated from labeled probe and detected by methods well-known to those of skill in the art, depending on the nature of the label. For example, radioactively labeled fragments can be separated by thin-layer chromatography and detected by autoradiography; while fluorescently-labeled fragments can be detected by irradiation at the appropriate excitation wavelengths with observation at the appropriate emission wavelengths. See, *e.g.*,
10 U.S. Patent No. 5,210,015.

In a variation of this technique, a probe contains both a fluorescent label and a quenching agent, which quenches the fluorescence emission of the fluorescent label. In this case, the fluorescent label is not detectable until its spatial relationship to the quenching agent has been altered, for example by exonucleolytic release of the fluorescent label from the
15 probe. Thus, prior to hybridization to its target sequence, the dual fluorophore/quencher labeled probe does not emit fluorescence. Subsequent to hybridization of the fluorophore/quencher-labeled probe to its target, it becomes a substrate for the exonucleolytic activity of a polymerizing enzyme which has initiated polymerization at an upstream primer. Exonucleolytic degradation of the probe releases the fluorescent label from the probe, and
20 hence from the vicinity of the quenching agent, allowing detection of a fluorescent signal upon irradiation at the appropriate excitation wavelengths. This method has the advantage that released label does not have to be separated from intact probe. Multiplex approaches utilize multiple probes, each of which is complementary to a different target sequence and carries a distinguishable label, allowing the assay of several target sequences simultaneously.

25 The use of MGB-modified oligonucleotide conjugates in this and related methods allows greater speed, sensitivity and discriminatory power to be applied to these assays. In particular, the enhanced ability of MGB-modified oligonucleotide conjugates to allow discrimination between a perfect hybrid and a hybrid containing a single-base mismatch will facilitate the use of hydrolyzable probe assays in the identification of single-
30 nucleotide polymorphisms and the like. One of skill in the art will appreciate that compositions and methods, such as those of the invention, that are capable of discriminating single-nucleotide mismatches will also be capable of discriminating between sequences that have 2, 3, 4, 5, or even 6 or more mismatches with respect to one another.

008277 6564260

In yet another related aspect, the present invention provides a method for primer extension, the method comprising incubating a polynucleotide containing a target sequence with one or more oligonucleotide primers complementary to the target sequence, in the presence of a polymerizing enzyme and nucleotide substrates under conditions favorable for polymerization; wherein at least one of the oligonucleotide primers contains a 3-substituted pyrazolo[3,4-*d*]pyrimidine or a 5-substituted pyrimidine base in place of a purine or pyrimidine base.

Amplification procedures are those in which many copies of a target nucleic acid sequence are generated, usually in an exponential fashion, by sequential polymerization and/or ligation reactions. In addition to the more traditional amplification reactions discussed below, the present invention is useful in amplifications involving three-way junctures (see, WO 99/37085), signal amplification (see Capaldi, et al., *Nuc. Acids Res.*, **28**:E21 (2000)), T7 polymerases, reverse transcriptase, RNase H, RT-PCR, Rolling Circles, cleavase and the like.

Many amplification reactions, such as PCR, utilize reiterative primer-dependent polymerization reactions. A primer is a nucleic acid that is capable of hybridizing to a second, template nucleic acid and that, once hybridized, is capable of being extended by a polymerizing enzyme (in the presence of nucleotide substrates), using the second nucleic acid as a template. Polymerizing enzymes include, but are not limited to, DNA and RNA polymerases and reverse transcriptases, *etc.* Conditions favorable for polymerization by different polymerizing enzymes are well-known to those of skill in the art. *See*, for example, Sambrook *et al.*, *supra*; Ausubel, *et al.*, *supra*; Innis *et al.*, *supra*. Generally, in order to be extendible by a polymerizing enzyme, a primer must have an unblocked 3'-end, preferably a free 3' hydroxyl group. The product of an amplification reaction is an extended primer, wherein the primer has been extended by a polymerizing enzyme.

Thus, in one embodiment of the invention, the methods and compositions disclosed and claimed herein are useful in improved amplification reactions such as PCR. *See, e.g.*, U.S. Patents 4,683,202; 4,683,195 and 4,800,159; Mullis and Faloona, *supra*; and Saiki *et al.*, *supra*. The polymerization step of PCR is most often catalyzed by a thermostable polymerizing enzyme, such as a DNA polymerase isolated from a thermophilic bacterium, because of the elevated temperatures required for the denaturation step of PCR. As discussed *supra*, one of the problems heretofore associated with the practice of PCR is the requirement for relatively long oligonucleotide primers, having sufficient hybrid stability to serve as primers at the elevated temperatures under which PCR is conducted. Modified oligonucleotides and especially MGB-modified oligonucleotide conjugates are useful as

primers in amplification reactions such as PCR, as the modified bases and MGBs increase hybrid stability, thereby significantly extending the lower limit of useful primer length. In addition, MGB-modified oligonucleotide conjugates are useful in specialized PCR protocols wherein reduced primer length is desirable. These include, but are not limited to, differential display, in which optimal primer length is below 10 nucleotides, random amplification of polymorphism in DNA (RAPD) techniques, and amplification length polymorphism analyses. Liang *et al, supra*; Williams *et al., supra*.

The modified oligonucleotides of the present invention are applicable to any type of assay or procedure in which PCR or a related amplification technique is used, including, but not limited to, hydrolyzable probe assays, priming with allele-specific oligonucleotides (ASOs), fragment length polymorphism analysis, single nucleotide polymorphism (SNP) analysis and microsatellite analysis, for example. These and other techniques are useful in gene mapping, in the identification and screening of disease-related genes, and in pharmacogenetics, to name just a few applications.

In still another related aspect, the present invention provides a method for determining the nucleotide sequence of a polynucleotide, the method comprising:

(a) incubating the polynucleotide with a modified oligonucleotide array under hybridization conditions; and

(b) determining to which of the modified oligonucleotides in the array the polynucleotide hybridizes;

wherein the modified oligonucleotides comprise at least one 3-substituted pyrazolo[3,4-*d*]pyrimidine in place of a purine base.

In these procedures, an ordered array comprising a plurality of modified oligonucleotides of different known sequences is used as a platform for hybridization to one or more test polynucleotides, nucleic acids or nucleic acid populations. Determination of the oligonucleotides which are hybridized and alignment of their known sequences allows reconstruction of the sequence of the test polynucleotide. See, for example, U.S. Patent Nos. 5,492,806; 5,525,464; 5,556,752; and PCT Publications WO 92/10588 and WO 96/17957. Materials for construction of arrays include, but are not limited to, nitrocellulose, glass, silicon wafers, optical fibers and other materials suitable for construction of arrays such as are known to those of skill in the art.

A major problem with current array-based sequencing and analysis methods is that the different oligonucleotides in an array will each have a different T_m . Hence, it is difficult to determine the stringency conditions that will provide maximum sensitivity, while

retaining the ability to distinguish single-base mismatches. This is a particularly important consideration for most, if not all, applications of array technology. Use of modified oligonucleotides and/or MGB-modified oligonucleotide conjugates in array-based sequencing and analysis techniques provides a solution to this problem. Surprisingly, conjugation of a MGB to a modified oligonucleotide makes its T_m relatively independent of base composition. Thus, for a population of modified oligonucleotides and MGB-modified oligonucleotide conjugates of a given length, the T_m for a perfect hybrid falls within a relatively narrow temperature range regardless of sequence. At the same time, the T_m for a single nucleotide mismatch is well below the T_m of the perfect match. Thus, arrays designed such that all modified oligonucleotides are the same length and are optionally present as their MGB conjugates exhibit minimal variation in T_m among the different oligonucleotides in the array, enabling more uniform hybridization conditions for the entire array. A further advantage to the use of modified oligonucleotides and MGB-modified oligonucleotide conjugates in these techniques is that it provides greater sensitivity, by allowing the use of shorter oligonucleotides, at higher temperatures (and hence higher stringency), while retaining single-nucleotide resolution.

An additional application of the present invention to array technology is in the examination of patterns of gene expression in a particular cell or tissue (see, generally, Eisen, et al., *METHODS IN ENZYMOLOGY*, **303**:179-205 (1999)). In this case, modified oligonucleotides or polynucleotides corresponding to different genes are arrayed on a surface, and a nucleic acid sample from a particular cell or tissue type, for example, is incubated with the array under hybridization conditions. Detection of the sites on the array at which hybridization occurs allows one to determine which modified oligonucleotides have hybridized, and hence which genes are active in the particular cell or tissue from which the sample was derived.

Array methods can also be used for identification of mutations or polymorphisms, where wild-type and mutant sequences are placed in an ordered array on a surface (see, Hacia, et al., *J. Mol. Genet.*, **36**:730-736 (1999)). Hybridization of a polynucleotide sample to the array under stringent conditions, and determination of which oligonucleotides in the array hybridize to the polynucleotide, allows determination of whether the polynucleotide possesses the wild-type or the mutant sequence. The increased discriminatory abilities of MGB-oligonucleotide conjugates are especially useful in this application of array technology.

Accordingly, the present invention provides a method for identifying a mutation in a target sequence of a gene of interest, the method comprising:

(a) incubating a polynucleotide comprising the target sequence with an array of oligonucleotides of different sequences, wherein the different sequences include the wild-type target sequence and different mutant target sequences, under hybridization conditions; and

(b) determining which of the oligonucleotides in the array become hybridized to the polynucleotide;

wherein one or more purine residues in a plurality of the oligonucleotides are replaced with a 3-substituted pyrazolo[3,4-*d*]pyrimidine.

In yet another related aspect, the present invention provides a method determining the nucleotide sequence of a target sequence in a polynucleotide, the method comprising:

(a) contacting a polynucleotide comprising the target sequence with at least two oligonucleotides of known sequence wherein one or more purine residues of the oligonucleotides are replaced by a 3-substituted pyrazolo[3,4-*d*]pyrimidine, and wherein one of the at least two oligonucleotides has a sequence that is perfectly complementary to the target sequence and at least one other of the oligonucleotides has a related target sequence and incubating each of the oligonucleotides with the polynucleotide under hybridization conditions; and

(b) determining the degree of hybridization between each of the oligonucleotides and the polynucleotide.

In one embodiment, a collection of all possible n-mer oligonucleotides (where n is an integer less than about 10) are used in a hydrolyzable probe assay to determine a nucleotide sequence. Each oligonucleotide is uniquely labeled (and preferably modified) and analysis of released label indicates which of the oligonucleotides has hybridized to the target sequence. Alignment of the sequences of the oligonucleotides which have hybridized provides the nucleotide sequence.

Modified oligonucleotides, and more preferably modified oligonucleotide-MGB conjugates are also useful in primer-dependent methods of DNA sequencing, such as the chain-termination method and its derivatives, originally described by Sanger *et al.*, *supra*. Use of MGB-modified oligonucleotide conjugates in chain-termination sequencing allows the use of shorter primers at higher stringency, and enables a greater degree of mismatch discrimination during sequencing. Examples include, but are not limited to, a search for

genes sharing a short region of homology (on the order of a few amino acids) and sequencing in a region in which very little existing sequence information is available. MGB-oligonucleotide conjugates are useful in such short primer sequencing techniques.

In still another embodiment, the present invention provides a method for
5 examining gene expression in a cell, the method comprising:

(a) incubating a population of polynucleotides representative of the genes expressed in the cell with an oligonucleotide array comprising a plurality of modified oligonucleotides of different sequences under hybridization conditions, and

(b) determining which of the modified oligonucleotides in the array become
10 hybridized to polynucleotides;


wherein said modified oligonucleotides comprise at least one 3-substituted pyrazolo[3,4-*d*]pyrimidine in place of a purine.

In one group of embodiments, the method is carried out with a plurality of expressed genes from multiple cell-types or tissues. The genes are preferably labeled with
15 different detection labels, then hybridized to an array at the same time and monitored via the label to determine each gene's expression pattern.


In addition to the assays and diagnostic methods described above, the modified oligonucleotides and modified bases described herein will have utility in antisense technologies. Antisense oligonucleotides are known to selectively inhibit gene expression and provide a genetic approach for disease treatment and prevention (Smith et al, *Int. J. Oncol.*, **17**:841-850 (2000)). Other modified oligonucleotides have been used to improve the performance of antisense oligonucleotides (Zhang et al, *Nat. Biotechnol.*, **18**:862-867 (2000); Flanagan et al, *Nat. Biotechnol.*, **14**:1139-1145 (1996)). Accordingly, another
20 embodiment of the present invention is the use of the 5-substituted pyrimidines, unsubstituted pyrazolo[3,4-*d*]pyrimidines and 3-substituted pyrazolo[3,4-*d*]pyrimidine as monomers, either
25 alone or in any combination, in the synthesis of antisense oligomers. In another embodiment the modified bases described herein can be used as monomers in an oligonucleotide to decrease enzymatic degradation of the antisense oligomers.

New Modified Bases

In another aspect, the present invention provides a number of new modified bases. These bases have the general formula:




wherein Z^1 is a member selected from the group consisting of H, F and OR^a wherein R^a is a member selected from the group consisting of H, (C_1-C_8) alkyl and a hydroxy protecting group; Z^2 is a member selected from the group of H, (C_1-C_8) alkyl, or is optionally combined with Z^1 to form a five- to seven-membered ring, having from one to three heteroatoms selected from the group consisting of O, S and N; Y^1 is a member selected from the group consisting of OH, a protected hydroxy group and $O-P^1$, wherein P^1 is a phosphoramidite or H-phosphonate group; Y^2 is a member selected from the group consisting of OH, a protected hydroxy group and $O-P^2$, wherein P^2 is a phosphoramidite, H-phosphonate, monophosphate, diphosphate or triphosphate; and B is a modified nucleotide selected from the group consisting of:




wherein X^{11} and X^{12} are each independently selected from the group consisting of H, NH_2 and a protected amino group; each R^{11} is independently selected from the group consisting of protected or unprotected forms of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl; and each R^{12} is independently selected from the group consisting of heterocyclyl(C_1-C_{12})alkyl, heterocyclyl(C_2-C_{12})alkenyl, heterocyclyl(C_2-C_{12})alkynyl, heterocyclyl, 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl, with the proviso that R^{12} is other than 2-pyridylethynyl.

In one group of embodiments, B is




Preferably, X^{11} is NH_2 and X^{12} is H. More preferably, X^{11} is NH_2 , X^{12} is H, Y^1 is $O-P^1$, Y^2 is a protected hydroxy, Z^1 is H, and R^{12} is selected from 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, 3-methoxyprop-1-ynyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 1,3-isoxazol-4-yl, 1,3-isoxazol-5-yl, 1,3-isoxazol-2-yl, 2-imidazolyl, 4-imidazolyl, 2-pyridyl, 3-pyridyl and 4-pyridyl. In particularly preferred embodiments, Y^1 is $-O-[(2\text{-cyanoethyl})\text{ N,N-diisopropylphosphoramidite}]$ and Y^2 is $-O-(4,4'\text{-dimethoxytrityl})$.

In another group of embodiments, B is



Preferably, X^{12} is NH_2 or H. More preferably, X^{12} is NH_2 or H, Y^1 is $O-P^1$, Y^2 is a protected hydroxy, Z^1 is H, and R^{12} is selected from 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, 3-methoxyprop-1-ynyl, 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 1,3-isoxazol-4-yl, 1,3-isoxazol-5-yl, 1,3-isoxazol-2-yl, 2-imidazolyl, 4-imidazolyl, 2-pyridyl, 3-pyridyl and 4-pyridyl. In particularly preferred embodiments, Y^1 is $-O-[(2\text{-cyanoethyl})\text{ N,N-diisopropylphosphoramidite}]$ and Y^2 is $-O-(4,4'\text{-dimethoxytrityl})$.



In still other preferred embodiments, the modified oligonucleotides of the invention have at least one base of formula III (including those in which the acetyl protecting groups are removed, and including other protected forms and activated forms thereof).



III


Preparation of compounds of formula III (including compounds wherein the 6-amino group is protected and the hydroxy groups of the sugar are either protected or activated as a phosphoramidite) are provided in Reaction Scheme 6, above.

In the most preferred embodiments of the invention, the modified bases are selected from formulae IVa, IVb or IVc.



These compounds are particularly well-suited for use in automatic
5 oligonucleotide synthesizers, and in preparing certain modified oligonucleotides described herein.


Other modified bases that are useful in the present invention include those represented by formula V:



V


wherein R^7 is selected from OH, SH or NH_2 . These compounds can be synthesized by methods described herein.

In another group of embodiments, B is




Preferably, X^{11} is NH_2 . More preferably, X^{11} is NH_2 , Y^1 is $O-P^1$, Y^2 is a protected hydroxy, Z^1 is H, and R^{11} is selected from 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, and 3-methoxyprop-1-ynyl. In particularly preferred embodiments, Y^1 is $-O-[(2\text{-cyanoethyl}) N,N\text{-diisopropylphosphoramidite}]$ and Y^2 is $-O-(4,4'\text{-dimethoxytrityl})$.

In another group of embodiments, B is




Preferably, Y^1 is $O-P^1$, Y^2 is a protected hydroxy, Z^1 is H, and R^{11} is selected from 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, and 3-methoxyprop-1-ynyl. In particularly preferred embodiments, Y^1 is $-O-[(2\text{-cyanoethyl}) N,N\text{-diisopropylphosphoramidite}]$ and Y^2 is $-O-(4,4'\text{-dimethoxytrityl})$.

In a related aspect, the present invention provides modified oligonucleotides having the formula:



wherein each Z^1 is independently selected from H, F and OR^a wherein R^a is a member selected from H, (C_1-C_8) alkyl and a hydroxy protecting group; each Z^2 is H or (C_1-C_8) alkyl, or is optionally combined with Z^1 to form a five- to seven-membered ring; each Z^3 is selected from O, S or NH; each Y is independently selected from $P(O)OH$, $P(S)OH$ and $P(O)CH_3$; the subscript n is an integer of from 1 to 98; W^1 and W^2 are each independently selected from H, a monophosphate, a diphosphate, a triphosphate and a minor groove binder-linking group moiety having an optionally attached reporter group or quencher; and each B is independently selected from the group consisting of adenine, guanine, cytosine, uridine and modified bases of the formula:



wherein X^{11} and X^{12} are each independently selected from H, NH_2 and a protected amino group; each R^{11} is independently selected from protected or unprotected forms of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl; and each R^{12} is independently selected from the group consisting of protected or unprotected forms of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, heterocyclyl (C_1-C_{12}) alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl and heterocyclyl, with the proviso that R^{12} is other than 2-pyridylethynyl; and

with the further proviso that at least one of the Bs is selected from the modified bases, and optionally, one or more of the Bs has an attached minor groove binder-linking group moiety, reporter group or a combination thereof.

Particularly preferred modified bases are those that have been described above.

EXAMPLES

In the hydrolyzable probe assay, a labeled probe is added to a PCR reaction. The probe is complementary to a region between the two PCR primers and is labeled with two fluorophores, one of which quenches the fluorescence of the other. The probe is designed to hybridize to its complementary target sequence on one of the PCR product strands at or above the strand extension temperature typically used in PCR ($55-75^\circ C$). The polymerizing enzymes normally used in PCR (*Taq* polymerase in particular) possess an

intrinsic 5'-exonuclease activity. During synthesis of new strands in the extension stage of the PCR reaction, this 5'-exonuclease activity will act on complementary strands bound to the template. If a probe, labeled as described above, is bound to the template, the 5'-exonuclease activity associated with the polymerizing enzyme will liberate the bound fluorophore. Once liberated, its fluorescence will no longer be quenched, and a fluorescent signal will be obtained. See, for example, U.S. Patent No. 5,210,015; Livak *et al.* (1995) *PCR Meth. App.* 4:357-362; and Heid *et al.* (1996) *Genome Res.* 6:986-994.

Thin-layer chromatography was run on silica gel 60 F-254 (EM Reagents) aluminum-backed plates. ¹H NMR were obtained at 300 MHz on a Varian VXR-300 spectrometer. Two-dimensional (Cosy) and NOE experiments assisted in the assignment of proton resonances. Elemental analyses were performed by Quantitative Technologies Inc. (Boundbrook, NJ).

Example 1

This example illustrates the synthesis of 5-(Prop-2-ynyl-4-methylbenzoate)-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxyuridine 3'-[(2-cyanoethyl) *N,N*-diisopropylphosphoramidite] (**3**).

5-(Prop-2-ynyl-4-methylbenzoate)-2'-deoxyuridine (1).

To a mixture of 5-iodo-2'-deoxyuridine (5.0 g, 14.12 mmol), CuI (270 mg, 1.42 mmol), Pd(PPh₃)₄ (0.82 g, 0.714 mmol), and triethylamine (2.4 mL) in 30 mL of anhydrous DMF was added prop-2-ynyl 4-methylbenzoate (6.10 g, 35.06 mmol). The mixture was stirred under argon for 4 h. and then evaporated to dryness. The residue was triturated in methanol and the excess prop-2-ynyl 4-methylbenzoate that precipitated was filtered away. The filtrate was evaporated and the residue was purified by silica gel chromatography eluting with 10% methanol in ethyl acetate. The pure product fractions were evaporated to dryness and the residue was precipitated from ethyl acetate- ether: 3.14 g (56%) yield; TLC (10% methanol in ethyl acetate), R_f = 0.50; ¹H NMR (DMSO-d₆) δ 11.68 (1H, s, uracil N-H), 8.29 (1H, s, 6-H), 7.89 and 7.35 (4H, 2 x d, J = 8.2 Hz, toluoyl Hs), 6.10 (1H, t, J = 6.4 Hz, 1'-H), 5.24 (1H, d, J = 4.4 Hz, 3'-OH), 5.15 (2H, s, alkynyl-CH₂), 5.12 (1H, t, J = 5.2 Hz, 5'-OH), 4.24 (1H, m, 3'-H), 3.79 (1H, q, 4'-H), 3.59 (2H, m, 5'-Hs), 2.39 (3H, s, toluoyl-CH₃), 2.13 (2H, m, 2'-H).

5-(Prop-2-ynyl-4-methylbenzoate)-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxyuridine (2).

To a solution of **1** (3.0 g, 7.50 mmol) in 45 mL of anhydrous pyridine was added dimethoxytrityl chloride (3.0 g). The resulting solution was stirred for 4 h. at room temperature and then poured into 400 mL of 5% sodium bicarbonate solution. The mixture was extracted with ethyl acetate (2 x 300 mL) and the extract was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with 5% methanol in the ethyl acetate. The pure product fractions were pooled and evaporated affording **2** as a foam: 4.16 g (79%) yield

5-(Prop-2-ynyl-4-methylbenzoate)-5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxyuridine 3'-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] (**3**).

To a solution of **2** (4.0 g, 5.70 mmol) in 130 mL of anhydrous methylene chloride, containing 3.0 mL of N,N-diisopropylethylamine was added chloro(2-cyanoethoxy)(N,N-diisopropylamino)phosphine (2.22 mL) under argon. The solution was stirred for 30 min at room temperature and then treated with 3.0 mL of methanol. The solution was diluted with 300 mL of ethyl acetate and washed with 300 mL of 5% sodium bicarbonate solution. The aqueous phase was extracted with 300 mL of ethyl acetate and the combined organic phases were dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of 30 to 0% hexane in ethyl acetate (2% triethylamine). The pure fractions were pooled and evaporated and the residue was precipitated from ethyl acetate – hexanes: 3.35 g (65%) yield; TLC (20% hexane in ethyl acetate), R_f = 0.82 and 0.71 (diastereomers); ³¹P NMR (DMSO-d₆) δ 147.82 and 147.45.

Example 2

Phase 1 Preparation of (2R,5R)-5-(4-amino-3-iodopyrazolo[3,4-d]pyrimidinyl)-2-(hydroxymethyl)oxolan-3-ol (**9**).

3-Iodo-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-one (**4**).

The synthesis of this compound was previously reported by Taylor et. al (*Tetrahedron*, **48(37)**:8089-8100 (1992)) using a N-iodosuccinimide. Our synthesis, employing iodine monochloride as the iodinating agent is described below.

To a 1.0 L solution of 1.2 M sodium acetate was added 4-hydroxypyrazolo[3,4-d]pyrimidine (25 g, 184 mmol) followed by iodine monochloride (46 g, 284 mmol). The mixture was stirred in a sealed container for 4 h. at 110 °C. The completed reaction was cooled to room temperature and treated with a solution of 30 g of sodium metabisulfite in 200 mL of water. The white precipitate that formed was filtered and rinsed with cold water. The solid was then dissolved in 200 mL of 3.2 M potassium hydroxide solution. Hydrazine monohydrate (5 mL) was added and the solution was stirred for 15 min. The solution was then acidified to ca. pH 7 by addition of 50 mL of concentrated HCl followed by a fine pH adjustment with acetic acid. The solid that formed was filtered, rinsed with cold water and dried: 55.1 g (114% yield). Note- it is difficult to remove residual water by drying the solid under vacuum. The residual water will be removed by a pyridine evaporation process in the next step.

3-Iodo-1,5-dihydropyrazolo[3,4-d]pyrimidin-4-thione (5).

Compound **4** (21.5 g, 82.1 mmol) was suspended in 150 mL of anhydrous pyridine and the mixture was evaporated to dryness. The solid was re-suspended in 170 mL of dry pyridine (under argon) and P₂S₅ (26.8 g) was added. The mixture was stirred for 10 min at 65 °C and then for 2-6 hours at 95 °C. The resulting solution was cooled to room temperature and degassed by bubbling a stream of argon into the solution (sulfide gas is passed into a trap containing sodium hydroxide solution). The reaction solution was then reduced in volume until a thick syrup formed. Excess P₂S₅ was decomposed by addition of 1 mL aliquots of water, until the vigorous reaction ceased, followed by addition of 500 mL of water and 10 mL of acetic acid. The mixture was warmed to 70 °C for 1 h to expedite expulsion of hydrogen sulfide gas and then diluted with 500 mL of water and cooled in an ice-bath. The solid was filtered, washed with water and dried: 19.8 g (87%) yield.

4-Ethylthio-3-iodopyrazolo[3,4-d]pyrimidine (6).

Compound **5** (43.5 g, 157 mmol) was stirred in potassium hydroxide solution (38.6 g KOH in 350 mL of water) for 30 min. The mixture was filtered and the filtrate was acidified to pH 10 by addition of acetic acid and then diluted with 350 mL of absolute ethanol. Iodoethane (10 mL) was added and the solution was stirred at room temperature.

Additional 10 mL aliquots of iodoethane were added after 30 min. and 1.0 h. The reaction was complete after a total of 90 min. Other runs required further addition of iodoethane to complete the reaction. The reaction solution was diluted with 700 mL of water and 20 mL of acetic acid. The mixture was cooled in an ice-bath and the solid was filtered, rinsed with water and dried. This crude product was dissolved in 300 of DMF and stirred at 90 °C for 15 min. Insoluble material was filtered away and the filtrate was diluted with 1 L of water and cooled in an ice-water bath. The solid was filtered, rinsed with water and dried: 19 g (40%) yield.

10 *[(2R,5R)-5-(4-Ethylthio-3-iodopyrazolo[3,4-d]pyrimidinyl)-3-(4-methylphenylcarbonyloxy)oxolan-2-yl]methyl 4-methylbenzonate (7).*

Compound 6 was converted to the corresponding potassium salt by reaction with 1 molar equivalent of potassium hydroxide in water. The resulting solution was evaporated to dryness and the residue was evaporated from dry acetonitrile. The potassium salt of 6 (29.0 g, 94.73 mmol) was dissolved in 80 mL of anhydrous DMF and then diluted with 830 mL of anhydrous acetonitrile. The chlorosugar derivative (48 g, 123 mmol) was added in one portion and the mixture was stirred for 3 h and then diluted with 1.5 L of 25% aqueous methanol. The mixture was allow to sit at 5 °C overnight. The crystals were filtered, rinsed with 25% aqueous methanol and dried: 32.2 g (52%) yield.

20 *(2R,5R)-2-(hydroxymethyl)-5-(3-iodo-4-methylpyrazolo[3,4-d]pyrimidinyl)oxolan-3-ol (8).*

To a suspension of 7 (10.97 g, 16.66 mmol) in 250 mL of methanol was added 22 mL of 1N sodium methoxide in methanol. The mixture was stirred at reflux and the reaction progress was monitored as soon as the mixture turned to a clear solution Note-formation of side products are observed if refluxing is continued after complete conversion to 5. The reaction was quenched by addition of 1.34 mL of acetic acid as soon as one component, corresponding to the desired product, is observed by TLC. The solution was evaporated and the residue was precipitated from methanol-ether-hexane. The solid was filtered and dried: 6.8 g (104%) yield crude product (contains sodium acetate).

30 *(2R,5R)-5-(4-amino-3-iodopyrazolo[3,4-d]pyrimidinyl)-2-(hydroxymethyl)oxolan-3-ol (9).*

Compound 8 (6.8 g, 17.35 mmol) was stirred in 200 mL of concentrated ammonium hydroxide (sealed container) at room temperature for 36 h. The mixture was

evaporated and the residue was precipitated from acetonitrile-ether. The solid was filtered and dried: 5.36 g (82%) yield. NMR and spectral data on this compound is identical to

Phase 2 Preparation of 3-{4-((1*E*)-1-aza-2-methylprop-1-enyl)-1-((2*R*,5*R*)-5-{[Bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-{[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}oxolan-2-yl)pyrazolo[3,4-*d*]pyrimidin-3-yl}prop-2-ynyl 4-methoxybenzoate (**13**; $R^1 = -OCOPhCH_3$)

3-{1-((2*R*,5*R*)-4-hydroxy-5-(hydroxymethyl)oxolan-2-yl)-4-aminopyrazolo[3,4-*d*]pyrimidin-3-yl}prop-2-ynyl 4-methylbenzoate (**10**; $R^1 = -OCOPhCH_3$)

To a mixture of **9** (2.40 g, 6.37 mmol), CuI (124 mg, 0.648 mmol), Pd(PPh₃)₄ (380 mg, 0.331 mmol), and triethylamine (1.32 mL) in 12 mL of anhydrous DMF was added prop-2-ynyl 4-methylbenzoate (1.87 g, 11.85 mmol). The mixture was stirred under argon for 12 h. and then evaporated to dryness. The residue was purified by silica gel chromatography eluting with 5% methanol in ethyl acetate. The pure product fractions were evaporated affording a foam: 2.29 g (85%) yield; TLC (10% methanol in ethyl acetate), $R_f = 0.43$; ¹H NMR (DMSO-*d*₆) δ 8.26 (1H, s, 6-H), 7.92 and 7.37 (4H, 2 x d, $J = 8.5$ Hz, toluoyl-Hs), 6.55 (1H, t, $J = 6.3$ Hz, 1'-H), 5.29 (2H, s, alkynyl-CH₂), 5.28 (1H, d, $J = 4.7$ Hz, 3'-OH), 4.76 (1H, t, $J = 5.7$ Hz, 5'-OH), 4.41 (1H, m, 3'-H), 3.81 (1H, m, 4'-H), 3.49 and 3.56 (2H, 2 x m, 5'-Hs), 2.76 and 2.24 (2H, 2 x m, 2'-Hs), 2.39 (3H, s, toluoyl-CH₃).

3-{4-((1*E*)-1-aza-2-methylprop-1-enyl)-1-((2*R*,5*R*)-4-hydroxy-5-(hydroxymethyl)oxolan-2-yl)pyrazolo[3,4-*d*]pyrimidin-3-yl}prop-2-ynyl 4-methylbenzoate (**11**; $R^1 = -OCOPhCH_3$).

Compound **10** (1.76 g, 4.16 mmol) was stirred in a solution of 5.0 mL of *N,N*-dimethylacetamide, 1.9 mL of *N,N*-dimethylacetamide dimethyl acetal and 2.0 mL of anhydrous triethylamine for 15 h. at room temperature. The solvents were evaporated and the residue was evaporated twice from xylenes affording **11** as a foam. TLC (10% methanol in ethyl acetate), $R_f = 0.29$; ¹H NMR (DMSO-*d*₆) δ 8.51 (1H, s, 6-H), 7.90 and 7.37 (4H, 2 x d, $J = 8.3$ Hz, toluoyl protons), 7.28, 7.14 and 6.74 (13H, d and 2 m, $J = 7.1$ Hz for the doublet), trityl protons), 6.63 (1H, m, 1'-H), 5.32 (1H, d, $J = 5.0$ Hz, 3'-OH), 5.24 (2H, s, alkynyl-CH₂), 4.56 (1H, m, 3'-H), 3.93 (1H, m, 4'-H), 3.68 (6H, s, methoxy Hs), 3.32 and 3.11 (6H, 2 x s, N-methyls), 3.09-2.95 (2H, m, 5'-Hs), 2.81 and 2.32 (2H, 2 x m, 2'-Hs), 2.40 (3H, s, -Me), 2.19 (3H, s, toluoyl-Me).

3-{4-((1E)-1-aza-2-methylprop-1-enyl)-1-((2R,5R)-5-{[Bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-hydroxyoxolan-2-yl)pyrazolo[3,4-d]primidin-3-yl]prop-2-ynyl 4-methylbenzoate (**12**; $R^1 = -OCOPhCH_3$)

To a solution of **11** in 25 mL of anhydrous pyridine was added dimethoxytrityl chloride (1.67 g). The resulting solution was stirred for 4 h. at room temperature and then poured into 250 mL of 5% sodium bicarbonate solution. The mixture was extracted with ethyl acetate and the extract was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with 5% methanol in the ethyl acetate. The pure product fractions were pooled and evaporated affording **12** as a foam: 2.06 g (62%) yield for the two-step process.

3-{4-((1E)-1-aza-2-methylprop-1-enyl)-1-((2R,5R)-5-{[Bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-{[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}oxolan-2-yl)pyrazolo[3,4-d]pyrimidin-3-yl]prop-2-ynyl 4-methoxybenzoate (**13**; $R^1 = -OCOPhCH_3$)

or,

1-[2-Deoxy-5-O-(4,4'-dimethoxytriphenylmethyl)- β -D-erythro-pentofuranosyl]-4-[(dimethylamino)prop-1-enyl]-3-(prop-2-ynyl-4-methylbenzoate) 1H-pyrazolo[3,4-d]pyrimidine 3'-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] (**13**).

To a solution of **12** (2.03 g, 2.56 mmol) in 60 mL of anhydrous methylene chloride, containing 1.33 mL of N,N-diisopropylethylamine was added chloro(2-cyanoethoxy)(N,N-diisopropylamino)phosphine (1.0 mL) under argon. The solution was stirred for 1.0 h at room temperature and then treated with 2.0 mL of methanol. The solution was diluted with 250 mL of ethyl acetate and washed with 200 mL of 5% sodium bicarbonate solution. The organic phase was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with 2% triethylamine in ethyl acetate. The pure fractions were pooled and evaporated. The phosphoramidite (**13**) was precipitated from ether-hexanes: 1.82 g (71%) yield; TLC (5 % methanol in ethyl acetate), $R_f = 0.32$; ^{31}P NMR (DMSO- d_6) δ 147.90 and 147.22.

Example 3

This example illustrates the preparation of N-{3-[1-((2R,5R)-5{[bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-[[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}oxolan-2-yl)-6-amino-4-oxo(5-hydropyrazolo[3,4-d]pyrimidin-3-yl)]propyl}-2,2,2-trifluoroacetamide (**22**)

3-Iodo-4-methoxypyrazolo[3,4-d]pyrimidine-6-ylamine(14).

4-Methoxy-pyrazolo[3,4-d]pyrimidin-6-ylamine (6.75 g, 40.87 mmol) was suspended in an aqueous solution containing sodium acetate (6.0 g, 44.09 mmol) and iodine monochloride (9.12 g, 56.17 mmol) and stirred at 100 °C in a sealed reaction vessel for 24 h. The resulting mixture was cooled to room temperature and treated with a solution of sodium metabisulfite (3.6 g, 18.94 mmol) in 24 mL of water. The solid that formed was filtered, rinsed with water and dried: 6.93 g (58%) yield; TLC (10% methanol in ethyl acetate), R_f = 0.57; ^1H NMR (DMSO- d_6) δ 13.08 (1H, br s, N-1 proton), 10.58 (1H, s, N-5 proton), 6.60 (2H, br s, 6-amino).

[(2R,5R)-5-(6-Amino-3-Iodo-4-methoxypyrazolo[3,4-d]pyrimidinyl)-3-(4-methylphenylcarbonyloxy)oxolan-2-yl]methyl 4-methylbenzoate (15).

To a suspension of (**14**) (6.68 g, 22.95 mmol) in 150 mL of methanol was added 8.05 mL of a 2.85 M methanolic potassium hydroxide. The mixture was stirred for one minute and then diluted with 100 mL of toluene and evaporated. The solid potassium salt of **1** was dried under vacuum. The potassium salt was suspended in 75 mL of anhydrous DMF and then diluted with 420 mL of anhydrous acetonitrile. 1-Chloro-1,2-dideoxy-3,5-di-O-toluoylribofuranose (8.95 g, 22.95 mmol) was immediately added and the reaction mixture was stirred at room temperature under argon for 1 h. and then filtered. The filtrate was evaporated and the residual foam was recrystallized twice from methanol: 6.59 g (45%) yield; TLC (1:1, ethyl acetate/hexane), R_f = 0.52; ^1H NMR (DMSO- d_6) δ 7.93 and 7.37 (8H, m and d, J = 7.9 Hz for the d, aromatic), 7.10 (2H, br s, 6-amino), 6.51 (1H, t, J = 6.6 Hz, 1'-H), 5.69 (1H, m, 3'-H), 4.44 (3H, m, 4'- and 5'-Hs), 3.99 (3H, s, 4-methoxy), 3.19 and 2.63 (2H, 2 x m, 2'-Hs), 2.40 and 2.38 (6H, 2 x s, toluoyl methyl protons).

[(2R,5R)-5-(6-Amino-3-Iodo-4-methoxypyrazolo[3,4-d]pyrimidinyl)-2-(hydroxymethyl)oxolan-3-ol (16).

To a suspension of **15** (32.4 g, 50.40 mmol) in 600 mL of methanol was added 12.5 mL of 1M sodium methoxide in methanol. The reaction mixture was stirred at reflux for 18 h. and then cooled in a freezer (-10 °C) . The crystals of **3** that formed were filtered and rinsed with ice-cold methanol: 10.95 g yield. Acetic acid (12.5 mL) was added to the filtrate and the volume was reduced to about 300 mL. The solution was allowed to sit in the freezer overnight and another crop of crystals (3.95 g) was collected. The filtrate was evaporated to dryness and the residual oil was triturated in ether. The solid that formed was filtered, dried and recrystallized from boiling water: 2.03 g. Total yield = 16.93 g (83%); TLC (10% methanol in ethyl acetate), R_f = 0.23; ^1H NMR (DMSO- d_6) δ 7.02 (2H, br s, 6-amino), 6.33 (1H, t, J = 6.1 Hz), 5.24 (1H, d, J = 4.4 Hz, 3'-OH), 4.73 (1H, t, J = 5.6 Hz, 5'-OH), 4.36 (1H, m, 3'-H), 3.99 (3H, s, 4-methoxy), 3.75 (1H, m, 4'-H), 3.51-3.32 (2H, m, 5'-Hs), 2.79 and 2.17 (2H, 2 x m, 2'-Hs). Anal. Calcd. For $\text{C}_{11}\text{H}_{14}\text{IN}_5\text{O}_4 \cdot 0.3 \text{ H}_2\text{O}$; C, 32.02; H, 3.57; N, 16.98. Found C, 32.13; H, 3.35; N, 16.77.

1-[(2R,5R)-(5-hydroxymethyl)oxolan-2-yl]-6-amino-3-iodo-5-hydropyrazolo[3,4-d]pyrimidin-4-one (17).

A suspension of **(16)** (16.84 g, 41.36 mmol) in 1.5 L of 1 N sodium hydroxide solution was heated to reflux. The mixture turned into a homogenous solution and was cooled to 25 °C using an ice-water bath. Acetic acid (90 mL) was added and the solution was stored at 5 °C overnight. The crystals that formed were filtered, rinsed with ice-cold water and dried: 7.47 g yield of **4**. The filtrate was evaporated to a volume of about 500 mL and stored at 5 °C overnight. A second crop of crystals was collected (3.75 g). The filtrate was evaporated to about 200 mL. The sodium acetate that formed was filtered off and the filtrate was stored at 5 °C overnight. Another crop of crystal was collected (2.25 g). Total yield: 13.47 g (83 %); TLC (5:3:2, n-butanol/water/acetic acid), R_f = 0.74; ^1H NMR (DMSO- d_6) δ 6.22 (1H, t, J = 6.3 Hz, 1'-H), 4.33 (1H, m, 3'-H), 3.73 (1H, m, 4'-H), 3.47 and 3.36 (2H, 2 x m, 5'-Hs), 2.63 and 2.12 (2H, 2 x m, 2'-Hs). Anal. Calcd. For $\text{C}_{10}\text{H}_{12}\text{IN}_5\text{O}_4 \cdot 0.7 \text{ H}_2\text{O}$; C, 29.60; H, 3.33; N, 17.26. Found C, 29.81; H, 3.02; N, 17.00.

N-(3-{1-[(2R,5R)-4-(hydroxymethyl)oxalan-2-yl]-6-amino-4-oxo(5-hydropyrazolo[3,4-d]pyrimidin-3-yl)}prop-2-ynyl)-2,2,2-trifluoroacetamide (18).

To a mixture of **17** (6.00 g, 15.26 mmol), CuI (297 mg, 1.56 mmol) and tetrakis[triphenylphosphine]palladium[0] in 30 mL of anhydrous DMF was added anhydrous triethylamine (3.14 mL) followed by propargyl trifluoroacetimide (4.29 g, 28.41 mmol). The reaction solution was stirred under argon for 40 h. The DMF was evaporated off and the residual oil was triturated in chloroform. The solid of crude **5** that formed was filtered, rinsed with chloroform and dried. The solid was dissolved in a minimum volume of DMF, absorbed onto silica gel and evaporated. The dry mixture was loaded onto a silica gel column and eluted with 10% methanol in ethyl acetate. The product fractions were pooled and evaporated. The residue was precipitated from ethyl acetate/ether: 4.0 g (63 %) yield; TLC (20% methanol in ethyl acetate), R_f = 0.59; ^1H NMR (DMSO- d_6) δ 10.79 (1H, s, $\text{N}_5\text{-H}$), 10.16 (1H, t, J = 5.2 Hz, trifluoroacetimido N-H), 6.77 (2H, br s, 6-amino), 6.28 (1H, t, J = 6.3 Hz, 1'-H), 5.23 (1H, d, J = 4.1 Hz, 3'-OH), 4.72 (1H, t, J = 5.1 Hz, 5'-OH), 4.32 (3H, m, -CONH- CH_2 - and 3'-H), 3.75 (1H, m, 4'-H), 3.50-3.29 (2H, 2 x m, 5'-Hs), 2.65 and 2.15 (2H, 2 x m, 2'-Hs). Anal. Calcd. For $\text{C}_{15}\text{H}_{15}\text{F}_3\text{N}_6\text{O}_5 \cdot 0.74 \text{H}_2\text{O}$; C, 41.93; H, 3.87; N, 19.56. Found C, 42.33; H, 3.64; N, 19.13.

N-(3-{1-[(2*R*,5*R*)-4-(hydroxymethyl)oxalan-2-yl]-6-amino-4-oxo(5-hydropyrazolo[5-4-*d*]pyriminin-3-yl)}propyl)-2,2,2-trifluoroacetamide (**19**).

To a solution of (**18**) (1.0 g, 2.40 mmol) in 100 mL of methanol, containing 0.12 g of 20% palladium hydroxide (preactivated with formic acid), was added 2.0 mL of 4 M triethylammonium formate buffer (pH 6.5). The mixture was shaken under 35 psi of hydrogen gas for 18 h (during some runs it was necessary to add additional catalyst for complete reduction). The mixture was filtered through Celite and the filtrate was evaporated. The residual oil was crystallized from water: 0.79 g (78%) yield; TLC (20% methanol in ethyl acetate), R_f = 0.52; ^1H NMR (DMSO- d_6) δ 10.59 (1H, s, $\text{N}_5\text{-H}$), 9.47 (1H, br t, trifluoroacetimido N-H), 6.64 (2H, br s, 6-amino), 6.27 (1H, t, J = 6.3 Hz, 1'-H), 5.18 (1H, d, J = 4.4 Hz, 3'-OH), 4.75 (1H, t, J = 5.9 Hz, 5'-OH), 4.36 (1H, m, 3'-H), 3.75 (1H, m, 4'-H), 3.48 and 3.61 (2H, 2 x m, 5'-Hs), 3.22, 2.68 and 1.87 (6H, 3 x m, propyl methylene protons), 2.68 and 2.12 (2H, 2 x m, 2'-Hs). Anal. Calcd. For $\text{C}_{15}\text{H}_{19}\text{F}_3\text{N}_6\text{O}_5 \cdot 0.90 \text{H}_2\text{O}$; C, 41.27; H, 4.80; N, 19.25. Found C, 41.57; H, 4.50; N, 19.11.

N-(3-[(1*E*)-1-aza-2-(dimethylamino)prop-1-enyl]-[(2*R*,5*R*)-4-(hydroxymethyl)oxalan-2-yl]-6-amino-4-oxo(5-hydropyrazolo[5-4-*d*]pyriminin-3-yl)}propyl)-2,2,2-trifluoroacetamide (**20**).

To a solution of **(19)** (0.80 g, 1.98 mmol) in 5.0 mL of anhydrous DMF was added N,N-dimethylformamide dimethylacetal (3.1 mL). The solution was stirred for 2.0 h. under argon and then evaporated. The residue was evaporated twice from xylenes and then placed under vacuum. The amorphous solid **(20)** that formed was triturated with ether and collected: 773-mg (82%) yield; TLC (20% methanol in ethyl acetate), $R_f = 0.47$; ^1H NMR (DMSO- d_6) δ 11.22 (1H, br s, $\text{N}_5\text{-H}$), 9.47 (1H, t, $J = 5.5$ Hz, trifluoroacetimido N-H), 8.67 (1H, s, $\text{N}=\text{CH-N}$), 6.42 (1H, t, $J = 6.5$ Hz, $1'\text{-H}$), 5.22 (1H, d, $J = 4.3$ Hz, $3'\text{-OH}$), 4.75 (1H, t, $J = 6.1$ Hz, $5'\text{-OH}$), 4.40 (1H, m, $3'\text{-H}$), 3.77 (1H, m, $4'\text{-H}$), 3.50 and 3.38 (2H, 2 x s, $5'\text{-Hs}$), 3.18 and 3.05 (6H, 2 x s, N,N-dimethyl Hs), 3.22, 2.72 and 1.89 (6H, 3 x m, propyl methylene Hs), 2.72 and 2.15 (2H, 2 x m, $2'\text{-Hs}$). Anal. Calcd. For $\text{C}_{18}\text{H}_{24}\text{F}_3\text{N}_7\text{O}_5 \cdot 0.40 \text{H}_2\text{O}$; C, 44.80; H, 5.18; N, 20.32. Found C, 45.02; H, 4.96; N, 19.94.

N-{3-[1-((2*R*,5*R*)-5-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-hydroxyoxolan-2-yl)-6-amino-4-oxo(5-hydropyrazolo[3,4-*d*]pyrimidin-3-yl)]propyl}-2,2,2-trifluoroacetamide (**21**).

To a solution of **(20)** (723-mg, 1.52 mmol) in 9.0 mL of anhydrous pyridine was added 4,4'-dimethoxytrityl chloride (0.61 g, 1.80 mmol). The reaction solution was stirred for 3.0 h. under argon and then poured into 100 mL of 5% sodium bicarbonate solution. The aqueous solution was extracted with ethyl acetate (2 x 200 mL) and the dried extracts (sodium sulfate) were evaporated. The crude product was purified by silica gel chromatography eluting with a solvent gradient of 0-5% methanol in ethyl acetate (2% triethylamine). The product fractions were evaporated affording **21** as an amorphous solid: 724-mg (61%) yield; TLC (5% methanol in ethyl acetate), $R_f = 0.39$; ^1H NMR (DMSO- d_6) δ 11.27 (1H, s, $\text{N}_5\text{-H}$), 9.43 (1H, t, $J = 5.3$ Hz, trifluoroacetimido N-H), 8.71 (1H, s, $\text{N}=\text{CH-N}$), 7.32, 7.17 and 6.76 (13H, 3 x m, aromatic), 6.45 (1H, t, $J = 6.3$ Hz, $1'\text{-H}$), 5.26 (1H, d, $J = 5.3$ Hz, $3'\text{-OH}$), 4.45 (1H, m, $3'\text{-H}$), 3.90 (1H, m, $4'\text{-H}$), 3.70 (6H, s, OMe Hs), 3.18 and 3.05 (10H, 2 x s, N,N-dimethyl, $5'\text{-Hs}$ and CONH-CH_2), 2.62 and 1.65 (4H, 2 x m, methylene Hs), 2.62 and 2.20 (2H, 2 x m, $2'\text{-Hs}$). Anal. Calcd. For $\text{C}_{39}\text{H}_{42}\text{F}_3\text{N}_7\text{O}_7 \cdot 0.30 \text{H}_2\text{O}$; C, 59.81; H, 5.48; N, 12.52. Found C, 59.80; H, 5.39; N, 12.63.

N-{3-[1-((2*R*,5*R*)-5-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}-4-{[bis(methylethyl)amino](2-cyanoethoxy)phosphinoxy}oxolan-2-yl)-6-amino-4-oxo(5-hydropyrazolo[3,4-*d*]pyrimidin-3-yl)]propyl}-2,2,2-trifluoroacetamide (**22**).

To a solution of (21) (700-mg, 0.900 mmol) in 22 mL of anhydrous methylene chloride, containing 0.47 mL of diisopropylethylamine, was added 2-cyanoethyl N,N-diisopropylchlorophosphoramidite (0.34 mL, 1.52 mmol). After stirring for 30 minutes under argon at 25 °C the solution was treated with 3.0 mL of methanol and diluted with 200 mL of ethyl acetate. The solution was washed with 100 mL of 5% sodium bicarbonate solution and dried over sodium sulfate and evaporated. The crude product was purified by silica gel chromatography eluting with 2% triethylamine in ethyl acetate. The product fractions were evaporated and the residue was precipitated from ether - hexane: 583-mg (66%) yield; TLC (ethyl acetate), R_f = 0.38; ^{31}P NMR (DMSO- d_6 , reference to 85% phosphoric acid) δ 145.50 and 144.72.

Example 4

This example illustrates the preparation of 3-{[5-(4,6-bis{(1E)-1-aza-2-[bis(2-methylpropyl)amino]vinyl}-3-prop-1-ynylpyrazolo[3,4-d]pyrimidinyl)-2-{[bis(4-methoxyphenyl)phenylmethoxy]-methyl}oxolan-3-yloxy][ethyl(methylethyl)amino]-phosphino}propanenitrile (26)

5-(4,6-bis{(1E)-1-aza-2-[bis(2-methylpropyl)amino]vinyl}-3-prop-1-ynylpyrazolo[3,4-d]pyrimidinyl)-2 (hydroxymethyl)oxolan-3-ol (24)

Compound (23) (1 mmol) can be stirred for 5 hr at room temperature with (dimethoxymethyl)bis(2-methylpropyl)amine (0.5 mL, 3.37 mmol). The reaction mixture can be reduced in vacuo, and the protected compound can be purified on silica gel to yield (24) (Vincent et al).

5-(4,6-bis{(1E)-1-aza-2-[bis(2-methylpropyl)amino]vinyl}-3-prop-1-ynylpyrazolo[3,4-d]pyrimidinyl)-2-{[bis(4-methoxyphenyl)phenylmethoxy]-methyl}oxolan-3-ol (25)

To a solution of (24) (1.50 mmol) in 9.0 mL of anhydrous pyridine 4,4'-dimethoxytrityl chloride (0.61 g, 1.80 mmol) can be added. The reaction solution can be stirred for 3.0 h. under argon and then can be poured into 100 mL of 5% sodium bicarbonate solution. The aqueous solution can be extracted with ethyl acetate (2 x 200 mL) and the extracts can be dried(sodium sulfate) and evaporated. The crude product can be purified by silica gel chromatography with an appropriate gradient to yield (25).

3-{[5-(4,6-bis{(1E)-1-aza-2-[bis(2-methylpropyl)amino]vinyl}-3-prop-1-ynyl)pyrazolo[3,4-d]pyrimidinyl)-2-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}-oxolan-3-yloxy][ethyl(methylethyl)amino}phosphino}propanenitrile (**26**)

To a solution of (**25**) (0.900 mmol) in 22 mL of anhydrous methylene chloride, containing 0.47 mL of diisopropylethylamine, 2-cyanoethyl N,N-diisopropylchlorophosphoramidite (0.34 mL, 1.52 mmol) can be added. The solution can be stirred for 30 minutes under argon at 25 °C and the can be treated with 3.0 mL of methanol and diluted with 200 mL of ethyl acetate. The solution can then be washed with 100 mL of 5% sodium bicarbonate solution and dried over sodium sulfate and evaporated. The crude product can be purified by silica gel chromatography with an appropriate gradient to yield (**26**).

Example 5

This example illustrates the synthesis of 5-[4,6-diamino-3-(2-methoxyethynyl)pyrazolo[3,4-d]pyrimidinyl]-2-(hydroxymethyl)oxolan-3-ol (**27**).

5-[4,6-diamino-3-(2-methoxyethynyl)pyrazolo[3,4-d]pyrimidinyl]-2-(hydroxymethyl)oxolan-3-ol (**27**)

To a mixture of 4,6-diamino-1-(2-deoxy-β-D-erythro-pentofuranosyl)-3-iodo-1H-pyrazolo[3,4-d]pyrimidine (4.2 g, 10.71 mmol), CuI (211 mg, 1.10 mmol), Pd(PPh₃)₄ (635 mg, 0.553 mmol), and triethylamine (2.2 mL) in 20 mL of anhydrous DMF was added methyl propargyl ether (1.82 mL). The mixture was stirred under argon for 16 h. and then evaporated to dryness. The residue was crystallized from methanol: 3.20 g (89%) yield; TLC (30% methanol in ethyl acetate), R_f = 0.63; ¹H NMR (DMSO-d₆) δ 7.44 (2H, br d, amino), 6.32 (1H, t, J = 6.6 Hz, 1'-H), 6.33 (2H, br s, amino), 5.20 (1H, d, J = 4.1 Hz, 3'-OH), 4.75 (1H, br t, 5'-OH), 4.40 (2H, s, methylene), 4.36 (1H, m, 3'-H), 3.76 (1H, m, 4'-H), 3.47 & 3.32 (2H, 2 x m, 5'-Hs), 3.32 (3H, s, methoxy), 2.68 & 2.14 (2H, 2 x m, 2'-Hs).

Synthesis of 5-{6-[(1E)-1-aza-2-(dimethylamino)vinyl]-4-[(1Z)-1-aza-2-(dimethylamino)vinyl]pyrazolo[3,4-d]pyrimidinyl}-2-(hydroxymethyl)oxolan-3-ol, 1-methoxyprop-1-yne (**28**).

Compound **27** (3.1 g, 9.28 mmol) was stirred in a solution of 30 mL of *N,N*-dimethylformamide and 15 mL of *N,N*-dimethylformamide dimethyl acetal for 14 h at 45 °C. The solvents were evaporated and the residue was evaporated twice from xylenes affording **28** as a foam which was precipitated from ethyl acetate - ether: 2.8 g (68%) yield; TLC (50% methanol in ethyl acetate), R_f = 0.36.

*Synthesis of 3-[(5-{6-[(1E)-1-aza-2-(dimethylamino)vinyl]-4-[(1Z)-1-aza-2-(dimethylamino)vinyl]pyrazolo[3,4-d]pyrimidinyl}-2-{[bis(4-methoxyphenyl)phenyl-methoxy]methyl}oxolan-3-yloxy)[bis(methylethyl)amino]phosphinooxy]propanenitrile (**29**).*

To a solution of **28** (2.7 g, 6.08 mmol) in 45 mL of anhydrous pyridine was added dimethoxytrityl chloride (2.4 g). The resulting solution was stirred for 3 h. at room temperature and then poured into 200 mL of 5% sodium bicarbonate solution. The mixture was extracted with ethyl acetate and the extract was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of 0-40% methanol in ethyl acetate. The pure 5'-*O*-DMT derivative fractions were pooled and evaporated affording a foam: 1.0 g (22%) yield.

To a solution of DMT derivative (0.98 g, 1.31 mmol) in 16 mL of anhydrous methylene chloride, containing 0.70 mL of *N,N*-diisopropylethylamine was added 2-cyanoethyl diisopropylchlorophosphoramidite (0.50 mL) under argon. The solution was stirred for 30 min at room temperature and then treated with 1.0 mL of methanol. The solution was loaded directly on to a silica gel column and eluted with a gradient of 0-20% methanol in ethyl acetate (2% triethylamine). The pure fractions were pooled and evaporated affording a foam: 0.25 g (20%) yield.

Example 6

This example illustrates the preparation of 3-[(5-{4-[(1Z)-1-aza-2-(methylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl}-2-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}oxolan-3-yloxy)[bis(methylethyl)amino]-phosphinooxy]propanenitrile (**35**) (See Reaction Scheme 8)

*(2-Furylmethoxymethylene)methane-1,1-dicarbonitrile (**30**)*

To an ice-cold solution containing anhydrous methylene chloride (500 mL), triethylamine (100 mL), and malonitrile (30g, 454 mmol) was added 2-furanyl chloride (50 g,

383 mmol) by dropwise addition over a 20 min period. Stirring was then continued for an additional hour at ambient temperature. The reaction solution was washed with 1.5 L of ice-cold 2N HCl solution and then with 1.5 L of water. The organic phase was evaporated affording an oil.

5 A portion of the product oil (4.6 g, 28.75 mmol) was dissolved in a solution consisting of 40 mL of dioxane and 4.0 mL of water. Dimethyl sulfate (15 mL) and sodium bicarbonate (15g) were added and the reaction solution was stirred for 2.5 h at 80 °C. The mixture was then dissolved in 100 mL of water and the product was extracted with 200 mL of ethyl acetate. The organic solution was washed with 100 mL of 5% sodium bicarbonate
10 solution followed by 100 mL of water. The solution was dried over sodium sulfate, filtered and evaporated affording an oil which solidified under vacuum: 2.72 g yield; TLC (1:1, ethyl acetate/hexane), R_f = 0.42; ^1H NMR ((DMSO- d_6) δ 8.25 (1H, m, aromatic), 7.63 (1H, m, aromatic), 6.92 (1H, m, aromatic), 4.23 (3H, s, methoxy).

15 *2-Amino-4-(2-furyl)furan-3-carbonitrile (31)* .

Hydrazine monohydrate (3.4 mL, 80 mmol) was added dropwise to an ice-cold solution of **30** (2.72 g, 15.63 mmol) in 75 mL of methanol over a period of 15 min. The solution was then evaporated to dryness affording an oil which solidified under vacuum. The solid was triturated in ether, filtered and dried: 2.2 g (81%) yield; TLC (ethyl acetate), R_f =
20 0.81; ^1H NMR (DMSO- d_6) δ 7.77 (1H, s, furanyl), 6.80 (1H, m, furanyl), 6.61 (1H, m, furanyl), 6.41 (2H, br s, amine).

3-(2-furyl)pyrazolo[3,4-d]pyrimidine-4-ylamine (32).

Compound **31** (25.7g, 148 mmol) was stirred in 250 mL of formamide at 190
25 °C for 4 h. The solution was then cooled to room temperature, diluted with 1.2 L of water and chilled in an ice-bath. The solid that formed was filtered and dried: 22 g (74%) yield; TLC (5% methanol in ethyl acetate), R_f = 0.25.

[5-(4-Amino-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl)-3-(4-methylphenylcarbonyloxy)-2-oxoethyl]oxolan-2-yl]methyl 4-methylbenzoate (33).

30 Compound **32** (10 g, 49.7 mmol) was stirred in 200 mL of 0.29 M methanolic KOH solution for 5 min. The mixture was evaporated to dryness and the residue was then dissolved in 40 mL of hot anhydrous DMF. The solution was cooled to room temperature

and then diluted with 230 mL of anhydrous acetonitrile. The chlorosugar derivative (23 g, 59.14 mmol) was added immediately and the mixture was stirred for 45 min and then evaporated to dryness. The residue was dissolved in 800 mL of ethyl acetate and washed with water (2 x 800 mL). The organic solution was dried over sodium sulfate, filtered and evaporated. The crude product was purified by silica gel chromatography eluting with a gradient of 30-0% hexane in ethyl acetate. The pure product fractions were evaporated and the residue was crystallized from hot methanol: 3.4 g (12%) yield; TLC (ethyl acetate), R_f = 0.47; ^1H NMR ((DMSO- d_6) δ 8.27 (1H, s, 6-H), 7.96 (1H, m, furanyl), 7.95, 7.84, 7.36 & 7.23 (8H, 4 x d, toluoyl aromatic), 6.83 (1H, m, furanyl), 6.80 (1H, t, J =6.3 Hz, 1'-H), 6.70 (1H, m, furanyl), 5.86 (1H, m, 3'-H), 4.64-4.42 (3H, m, 4'-H and 5'-Hs), 3.28 & 2.76 (2H, 2 x m, 2'-Hs), 2.39 & 2.36 (6H, 2 x s, toluoyl methyls).

5-{4-[(1Z)-1-aza-2-(dimethylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl}-2-(hydroxymethyl)oxolan-3-ol (34).

A suspension of **33** (3.36 g, 6.08 mmol) in 17 mL of 0.12 M methanolic sodium methoxide solution was refluxed for 30 min. The resulting solution was cooled to room temperature and neutralized by addition of 0.12 mL of acetic acid. The solution was evaporated to dryness and the product was precipitated from methanol-ether and then recrystallized from boiling water: 1.63 g (85%) yield of the deprotected nucleoside.

All of the nucleoside product from above was stirred in a solution consisting of 30 mL of anhydrous DMF and 15 mL of N,N-dimethylformamide dimethylacetal for 5 h. The solution was evaporated to dryness and the residue was then evaporated twice from xylenes affording a foam: 1.89 g (99%) yield; TLC (20% methanol in ethyl acetate), R_f = 0.45; ^1H NMR ((DMSO- d_6) δ 8.98 & 8.46 (2H, 2 x s, formamidine C-H and 6-H), 7.97 (1H, m, furanyl), 7.81 (1H, m, furanyl), 6.65 (2H, m, 1'-H and furanyl), 5.30 (1H, d, J = 4.5 Hz, 3'-OH), 4.79 (1H, t, J = 5.5 Hz, 5'-OH), 4.51 (1H, m, 3'-H), 3.85 (1H, m, 4'-H), 3.57 & 3.40 (2H, 2 x m, 5'-Hs), 3.26 & 3.21 (6H, 2 x s, N-Me protons), 2.88 & 2.30 (2H, 2 x m, 2'-Hs).

3-[(5-{4-[(1Z)-1-aza-2-(methylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl}-2-[[bis(4-methoxyphenyl)phenylmethoxy]methyl]oxolan-3-yloxy)[bis(methylethyl)amino]-phosphinoxy]propanenitrile (35).

Dimethoxytrityl chloride (2.01 g) was added to a solution of **34** (1.84 g, 4.95 mmol) dissolved in 30 mL of dry pyridine. The reaction solution was stirred for 3 h at room temperature and then poured into 200 mL of 5% sodium bicarbonate solution. The product was extracted with 300 mL of ethyl acetate and the organic phase was dried over sodium sulfate and evaporated. The residue was purified using silica gel chromatography eluting with a gradient of 0-5% methanol in ethyl acetate. The pure product fractions were evaporated affording a foam: 2.26 g (68%) yield of the 5'-*O*-DMT derivative.

To a solution of the DMT derivative (2.18 g, 3.23 mmol) dissolved in 40 mL of dry methylene chloride, containing 1.75 mL of *N,N*-diisopropylethylamine, was added 1.25 mL of 2-cyanoethyl diisopropylchlorophosphoramidite. The solution was stirred under argon for 15 min at room temperature and then treated with 5 mL of methanol. The resulting solution was diluted with 500 mL of ethyl acetate and washed with 400 mL of 5% sodium bicarbonate solution. The organic solution was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of 5-10 % methanol in ethyl acetate (2% triethylamine). The pure product fractions were evaporated affording a foam: 1.62 g (57%) yield; ³¹P NMR (DMSO-*d*₆) δ 147.81 and 147.16.

Example 7

This example illustrates the preparation of 3-[(5-{6-[(1*E*)-1-aza-2-(dimethylamino)vinyl]-4-[(1*Z*)-1-aza-2-(dimethylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-*d*]pyrimidinyl}-2-[[bis(4-methoxyphenyl)phenylmethoxy]methyl}oxolan-3-yloxy)[bis(methylethyl)amino]phosphinoxy]propanenitrile (**39**, See Reaction Scheme 9)

*3-(2-furyl)pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (36)*

A mixture of finely ground **31** (10.0 g, 57.47 mmol) and guanidine carbonate (16.6 g, 91.95 mmol) was heated at 230 °C for 45 min. The mixture was cooled to room temperature and the solid was triturated in 100 mL of boiling water. The solid (pure **36**) was filtered, rinsed with water and dried: 11.1 g (89%) yield; TLC (40% methanol in ethyl acetate), *R*_f = 0.66; ¹H NMR ((DMSO-*d*₆) δ 12.68 (1H, br s, N-H), 7.86 (1H, m, furanyl), 6.93 (2H, br s, -NH₂), 6.85 (1H, d, *J* = 3.5 Hz, furanyl), 6.65 (1H, m, furanyl), 6.10 (2H, br s, -NH₂).

5-(4,6-diamino-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl)-2-(hydroxymethyl)oxolan-3-ol (**37**)

Compound **36** (10.5 g, 48.61 mmol) was stirred in 200 mL of 0.29 M methanolic KOH solution for 5 min. The mixture was evaporated to dryness and the residue was then dissolved in 105 mL of hot anhydrous DMF. The solution was cooled to room temperature and then diluted with 620 mL of anhydrous acetonitrile. The chlorosugar derivative (23 g, 59.14 mmol) was added immediately and the mixture was stirred for 40 min and then filtered. The filtrate was evaporated to dryness and the residue was chromatographed through a silica gel column eluting with ethyl acetate. The nucleoside fractions were evaporated affording 2.8 g (10 % yield) of the toluoyl protected nucleoside as a mixture of alpha- and beta-anomers. This mixture was refluxed in 40 mL of 0.19 M methanolic sodium methoxide for 45 min. The reaction solution was placed in a freezer overnight producing a crop of crystals corresponding to the pure beta-anomer **37**: 690 mg yield; TLC (20% methanol in ethyl acetate), R_f = 0.32; ^1H NMR ($\text{DMSO}-d_6$) δ 7.90 (1H, m, furanyl), 6.99 (2H, br s, amino), 6.86 (1H, d, J = 4.1 Hz, furanyl), 6.68 (1H, m, furanyl), 6.41 (1H, t, J = 6.6 Hz, 1'-H), 6.26 (2H, br s, amino), 5.21 (1H, d, J = 4.4 Hz, 3'-OH), 4.82 (1H, t, J = 5.8 Hz, 5'-OH), 4.42 (1H, m, 3'-H), 3.79 (1H, m, 4'-H), 3.52 & 3.41 (2H, 2 x m, 5'-Hs), 2.75 & 2.18 (2H, 2 x m, 2'-Hs).

5-{6-[(1E)-1-aza-2-(dimethylamino)vinyl]-4-[(1Z)-1-aza-2-(dimethylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl}-2-(hydroxymethyl)oxolan-3-ol (**38**).

A solution of **37** (0.68 g, 2.05 mmol) in 30 mL of anhydrous DMF and 15 mL of N,N-dimethylformamide dimethylacetal was stirred at room temperature for 24 h. The solution was evaporated to dryness and the residue was evaporated twice from xylenes affording a foam: 0.90 g (99%) yield; TLC (50 % methanol in ethyl acetate), R_f = 0.38; ^1H NMR ($\text{DMSO}-d_6$) δ 8.93 & 8.77 (2H, 2 x s, formamidine C-Hs), 7.94 (1H, m, furanyl), 7.76 (1H, m, furanyl), 6.62 (1H, m, furanyl), 6.60 (1H, t, J = 6.6 Hz, 1'-H), 5.25 (1H, d, J = 4.4 Hz, 3'-OH), 4.84 (1H, t, J = 5.9 Hz, 5'-OH), 4.47 (1H, m, 3'-H), 3.83 (1H, m, 4'-H), 3.56 & 3.41 (2H, 2 x m, 5'-Hs), 3.25, 3.18, 3.16 & 3.03 (12H, 4 x s, N-methyls), 2.82 & 2.22 (2H, 2 x m, 2'-Hs).]

3-[(5-{6-[(1E)-1-aza-2-(dimethylamino)vinyl]-4-[(1Z)-1-aza-2-(dimethylamino)vinyl]-3-(2-furyl)pyrazolo[3,4-d]pyrimidinyl}-2-{[bis(4-methoxyphenyl)phenylmethoxy]methyl}oxolan-3-yloxy)[bis(methylethyl)amino]phosphinoxy]propanenitrile (**39**).

Dimethoxytrityl chloride (0.85 g) was added to a solution of **38** (0.90 g, 2.04 mmol) in 12 mL of dry pyridine. The reaction solution was stirred for 2 h at room temperature and then poured into 200 mL of 5% sodium bicarbonate solution. The product was extracted with ethyl acetate (3 x 300 mL) and the pooled organic extracts were dried over sodium sulfate and evaporated. The residue was purified using silica gel chromatography eluting with a gradient of 20-30% methanol in ethyl acetate. The pure product fractions were evaporated affording a foam: 286-mg yield of the 5'-O-DMT derivative.

To a solution of the DMT derivative (286 mg, 0.384 mmol) dissolved in 5 mL of dry methylene chloride, containing 0.23 mL of N,N-diisopropylethylamine, was added 0.17 mL of 2-cyanoethyl diisopropylchlorophosphoramidite. The solution was stirred under argon for 15 min at room temperature and then treated with 0.5 mL of methanol. The resulting solution was diluted with 100 mL of ethyl acetate and washed with 75 mL of 5% sodium bicarbonate solution. The organic solution was dried over sodium sulfate, filtered and evaporated. The residue was purified by silica gel chromatography eluting with a gradient of 0-30 % methanol in ethyl acetate (2% triethylamine). The pure product fractions were evaporated affording a foam: 230 mg (12 %) yield; ³¹P NMR (DMSO-*d*₆) δ 147.77 and 147.08.

Example 8

This example illustrates the enhanced ability of modified oligonucleotides to discriminate between related target sequences.

In this example, modified oligonucleotides were prepared containing PPPA, PPG and a minor groove binder. As Figure 3 illustrates, the use of a MGB in combination with PPPA and PPG increases the *T*_m and allows the design of shorter probes compatible with PCR extension temperatures. These modified oligonucleotides also yield increased mismatch discrimination. In a first design of a 18-mer fluorescein-ODN-Red 13quencher-MGB real-time PCR probe, poor discrimination of an A/T mismatch was observed as shown in Figure 3A. Re-design of the probe against the opposite strand, putting the mismatch under

the MGB and substituting PPPA for A and PPG for G as indicated, allows shortening of the probe to a 15-mer. This probe now gives good mismatch discrimination as shown in Figure 3B.

A thermodynamic investigation of mismatch discrimination was performed on a set of oligonucleotides hybridized to a set of targets perfectly match or containing a single mismatch. The target sequences contain a) normal As, b) PPPA, c) normal As and a 3'-MGB and d) PPPA and a 3'-MGB, respectively. The sequences for the probes and targets are shown below in Table 3a and 3b. The determination of T_m s and the calculation of

Table 3a and 3b
Probe and target sequences

A. Probe Sequences – Mismatch Underlined		
Number	Mismatch	Probe Sequence
1	Complement	AAAGTTATGTCTACTTACAGAAA
2	A/C	AAAGCTATGTCTACTTACAGAAA
3	A/C	AAAGTCATGTCTACTTACAGAAA
4	T/G	AAAGTTGTGTCTACTTACAGAAA
5	A/C	AAAGTTACGTCTACTTACAGAAA
6	C/A	AAAGTTATATCTACTTACAGAAA
7	A/C	AAAGTTATGCCTACTTACAGAAA
8	G/T	AAAGTTATGTTTACTTACAGAAA
9	A/C	AAAGTTATGTCCACTTACAGAAA
10	T/G	AAAGTTATGTCTGCTTACAGAAA
11	G/T	AAAGTTATGTCTATTACAGAAA
12	A/C	AAAGTTATGTCTACCTACAGAAA
13	A/C	AAAGTTATGTCTACTCACAGAAA
14	T/G	AAAGTTATGTCTACTTGCAGAAA
B. Target Sequences – A'=PPPA		
1		GTAAGTAGACATAAC
2		GTA'A'GTA'GA'CA'TA'A'C
3		GTAAGTAGACATAAC-MGB
4		GTA'A'GTA'GA'CA'TA'A'C-MGB

Table 3c
Thermodynamic comparison of mismatch discrimination in terms of free energy increment at 50°C where $\Delta\Delta G^{\circ}_{50} = R^{\circ} \ln(K_{\text{match}}/K_{\text{mismatch}})$

Number	Mismatch	A	PPPA	MGB	MGB+PPPA
2	A/C	2340	2930	2870	5320
3	A/C	2560	3280	4100	6320
4	T/G	1950	1810	4200	5900
5	A/C	3520	3760	3830	4980
6	C/A	5030	5340	4190	5970

7	A/C	3000	3370	4310	5260
8	G/T	3040	3260	3070	4820
9	A/C	3290	3440	3810	5630
10	T/G	1800	1950	2090	3350
11	G/T	3340	3120	3630	5070
12	A/C	2940	3620	2550	4490
13	A/C	2360	3210	1820	3980
14	T/G	1600	2010	2000	2480

$\Delta\Delta G_{50}^{\circ}$ is described in Example 9. Table 3c clearly shows increased mismatch discrimination when PPPA is substituted for A and even larger discrimination when PPPA is combined with a MGB.

- 5 Comparison of the thermodynamic discrimination of mismatched base pairs formed in MGB-ODNs containing HO-PPPA/HO-PU with PPPA/PU at 37°C is shown in Table 4. The ODNs containing the modified bases in combination with MGB were hybridized to their complements. Mismatches are underlined in the sequences shown in Table 4. As shown in this table, HO-PPPA and HO-PU substitution compared to PPPA and
- 10 PU, shown for the most part increased mismatch discrimination.

Table 4

Comparison of thermodynamic discrimination of mismatched base pairs formed by HOPPPA or HOPU vs PPPA and PU in the 8-mer duplexes (+MGB).

15

	Sequence of Duplex	PPPA/ PU $\Delta\Delta G$ cal/mol	HOPPPA/ HOPU $\Delta\Delta G$ cal/mol		Sequence of	PPPA/ PU $\Delta\Delta G$ cal/mol	HOPPPA/ HOPU $\Delta\Delta G$ cal/mol
Match	CGUCACUG-MGB AGCTGTGACT			Match	UAUU <u>A</u> UUG-MGB AATAATAACC		
1	CGU <u>C</u> ACUG-MGB AGCTGTGACT	4250	4350	10	UAU <u>U</u> AUUG-MGB AATTATAACC	4400	5000
2	CGU <u>C</u> ACUG-MGB AGC <u>G</u> GTGACT	3450	3540	11	UAU <u>U</u> AUUG-MGB AATGATAACC	3740	3760
3	CGU <u>C</u> ACUG-MGB AGC <u>C</u> GTGACT	4860	4530	12	UAU <u>U</u> AUUG-MGB AATCATAACC	6630	6840
4	CGUC <u>A</u> CUG-MGB AGCAG <u>A</u> GACT	4870	4850	13	UAUU <u>A</u> UUG-MGB AATA <u>A</u> AAACC	5090	5730
5	CGUC <u>A</u> CUG-MGB AGCAGG <u>G</u> GACT	4190	4360	14	UAUU <u>A</u> UUG-MGB AATAAG <u>A</u> AACC	5920	6520
6	CGUC <u>A</u> CUG-MGB AGCAGC <u>G</u> GACT	3930	3940	15	UAUU <u>A</u> UUG-MGB AATAAC <u>A</u> AACC	4120	4530
7	CGU <u>C</u> ACUG-MGB AGCA <u>A</u> TGACT	2600	2300				
8	CGU <u>C</u> ACUG-MGB AGCA <u>T</u> TGACT	4360	4210				
9	CGUC <u>A</u> CUG-MGB AGCACTGACT	4420	4610				

$\Delta\Delta G$ was calculated at 37°C.

Example 9

5 This example illustrates the UV melting studies carried out using oligonucleotides of the present invention.

Hybrids formed between unmodified ODNs or MGB-probes and their complements were melted at a rate of 0.5°C/min, on a Lambda 2S (Perkin-Elmer) spectrophotometer with a PTP-6 automatic multicell temperature programmer. T_m data was prepared using 0.5x SSPE buffer (Sigma, pH 7.4). Each ODN (1uM of each strand) was mixed with its complement to give a 1:1 ratio. Prior to melting, samples were denatured at 100°C and then cooled to 10°C over a 10 min period. Mismatch discrimination for each type of duplex was calculated in terms of $\Delta\Delta G$ at 50°C using the equation:

$$\Delta\Delta G_{50}^{\circ} = R^{\circ}\ln(K_{\text{match}}/K_{\text{mismatch}}).$$

15 The term $K_{\text{match}}/K_{\text{mismatch}}$ can be determined using the relative fractions of duplex and single strands calculated from the melting curves at 50°C (see Lohkov, S.G. & Pyshnyi, *FEBS Letters* **420**:134-138 (1997)).

Example 10

20 This example illustrates the hybridization of DNA to oligonucleotide glass microarrays prepared as described in co-pending application Ser. No. 09/364,320 and further optimized for 8-10 mer MGB probes.

25 DNA (oligonucleotide or PCR amplicon) at $1-5 \times 10^{-7}$ M in 5XSSPE, 0.1% Triton X-100, 10% formamide was hybridized to the microarray in Frame-Seal Chambers (MJ Research) under the following conditions: 5 minutes at 55°C, slow cooling at 0.1 °C/sec to 35 °C, 60 minutes at 55°C. The slides were then washed in 0.5XSSPE for 5 minutes at 45°C. Slides were dried under a stream of air and scanned using an Array Works Fluorescent scanner (Applied Precision). The washing procedure can be repeated under more stringent conditions if required.

30

Example 11

This example illustrates a single base primer extension assay.

5 Single stranded template DNA and 6-mer primer (20 μ M each), 1X Thermopol buffer (New England Biolabs), 10mM manganese chloride (USB), were combined and incubated 5 minutes at 50°C. Then 5 μ Ci α -³²P ddATP and 8 units Bst polymerase (NEB) were added, bringing total volume to 10 μ l, and incubated 15 minutes at 50°C. After
10 incubation, 6 μ l stop solution (95% formamide, 20mM EDTA) was added and reactions were cooled to room temperature. Samples were diluted 1:10 in denaturing dye (35% formamide, 0.05% xylene cyanol, 0.05% bromophenol blue, 1mM EDTA), and aliquots were analyzed by electrophoresis on a denaturing 10% polyacrylamide gel.

 The use of modified bases in primer extension assays for SNP typing is illustrated in Table 5. Primer extension with Bst polymerase (NEB) was evaluated with a
15 primer substituted with different modified bases and the amount of product was measured based on the incorporated radioactivity using gel electrophoresis.

Table 5

Comparison of ³²P-incorporation in primer extension product by polyacrylamide
20 gel electrophoresis using AAC CAC TCT GTC CTA template

Primer	Relative Signal
TGAGAC	++
U ^{prop} GAGAC	+
U ^{prop} G ^{prop} AGAC	+++
U ^{prop} G ^{prop} A ^{prop} GAC	+++++
U ^{prop} G ^{prop} A ^{prop} G ^{prop} AC	+++++
U ^{prop} G ^{prop} A ^{prop} G ^{prop} A ^{prop} C	+
U ^{prop} G ^{prop} A ^{prop} G ^{prop} A ^{prop} C ^{prop}	0.5+

Example 12

This example illustrates the use of the algorithm described in the specification to predict the T_m of modified oligonucleotides containing PPG both with and without a minor groove binder (CDPI₃)

Using thermodynamic parameters in Tables 2a/2b and the correction formula for the MGBs contribution, T_m of PPG oligonucleotides with and without a MGB can be calculated with algorithm procedures as described above. The thermodynamic parameters of the nearest neighbor pairs that are not G-containing do not change when the oligonucleotides are substituted with PPG. Similarly the thermodynamic parameters of the nearest-neighbor pairs do not change when MGB containing oligonucleotides are substituted with PPG.

Table 6 illustrates the ability of the algorithm to predict the T_m of oligonucleotides containing PPG alone or in combination with a minor groove binder (CDPI₃). In Table 6, all Gs are substituted with PPGs.

Table 6

Comparison of experimental T_m s with that of predicted T_m s using the nearest-neighbor thermodynamic parameters for PPG containing oligonucleotides and PPG containing oligonucleotides attached to a MGB

5'-Probe Sequence -3'	ODN Duplex Stability °C			MGB-ODN Duplex Stability °C		
	$T_{m_{exp}}$	$T_{m_{calc}}$	Err	$T_{m_{exp}}$	$T_{m_{calc}}$	Err
CTGTAAGTAGATATAAC	51.84	53.23	1.39	65.88	66.69	0.81
GGCAAGATATATAG	50.21	49.81	-0.40	66.37	65.56	-0.81
GTGACGCAGATTCC	61.27	61.06	-0.21	76.97	75.19	-1.78
GTAAGTAGACATAAC	52.12	51.78	-0.34	64.64	63.31	-1.33
CAGGGAGCTTTGGA	59.9	60.22	0.32	74.39	71.47	-2.92
CACTCGTGAAGCTG	60.85	59.49	-1.36	74.04	72.26	-1.78
GTAAGTAGGCATAAC	55.74	55.47	-0.27	66.91	66.00	-0.91
CCGGATGTAGGATC	57.52	59.05	1.53	69.3	70.03	0.73
GATTACCTGGATT	50.64	50.32	-0.32	62.29	62.33	0.04
CCGTCAATGGTCAC	58.66	60.01	1.35	70.13	69.91	-0.22
CAGCACGTAGCC	57.31	58.07	0.76	69.29	67.60	-1.69
CGGCTACGTGCTGG	65.19	66.01	0.82	76.12	74.79	-1.33
CGGCTACATGCTGG	61.14	61.95	0.81	71.56	72.99	1.43
CTAAATCTGCCG	50.4	48.09	-2.31	62.08	60.19	-1.89
TCTGGATGATGGGCA	61.74	61.95	0.21	71.65	72.13	0.48
GTTTCATGGGTGTAAT	57.51	57.77	0.26	66.94	68.79	1.85
CGGAGGTAGGATCA	59.24	59.46	0.22	69.46	70.93	1.47
CCACCCGCCTCAG	60.73	61.14	0.41	71.43	70.74	-0.69
CACAGGAGTGGTTGG	63.07	64.40	1.33	72.28	72.92	0.64

CGGACCAGTGCGTG	68.1	67.58	-0.52	77.92	76.80	-1.12
TCGGACCAGTGCGT	65.04	66.00	0.96	74.94	75.62	0.68
AACGGGGTACGATA	57.93	57.11	-0.82	67.79	67.08	-0.71
CAGTTGAGATTCTAAGAC	60.06	60.15	0.09	67.15	67.43	0.28
AGGGGCGTCTTG	60.78	58.57	-2.21	71.62	72.76	1.14
GTAAGTAGGCATAGC	58.34	58.95	0.61	65.95	66.99	1.04
TGCCCAGCCCCAG	63.13	63.40	0.27	71.28	71.32	0.04
CCAACACTCGTGAA	54.87	56.14	1.27	62.07	63.54	1.47
GTAAGTAGACACAGC	59.48	58.41	-1.07	65.79	66.27	0.48
TCGGACCAGTGC	58.02	58.55	0.53	65.99	66.35	0.36
CGATCACGCTGGC	62.12	62.75	0.63	69.18	71.81	2.63
GTCCTGGGGGTGG	65.19	64.54	-0.65	72.78	72.53	-0.25
GTAAGTAGGTGTGAC	60.7	59.70	-1.00	66.92	67.00	0.08
GGTTGTACGGGTTTACG	68.38	68.81	0.43	74.16	75.38	1.22
GGACCAGTGCGTGA	66.84	65.46	-1.38	73.38	71.53	-1.85
GTAAGTAGACGCAGC	62.91	62.44	-0.47	68	67.82	-0.18
GTAAGTAGGCGCAGC	65.52	65.91	0.39	69.8	70.34	0.54
GTAAGTAGGCGCGGC	68.71	68.96	0.25	72.26	72.76	0.50
GGTTCCCGAGCG	62.15	61.14	-1.01	65.75	64.22	-1.53

The accuracy of the prediction algorithm is about +/- 1 and +/- 2°C for PPG containing oligonucleotides and PPG containing MGB-oligonucleotides, respectively.

Accordingly, a sequence of interest can be obtained from a source such as Genbank, and a T_m window can then be set as a requirement for a set of probes or primers. Using the algorithm above, and information from the nearest-neighbor parameters, a collection of probe or primer sequences with the desired T_m s can be calculated.

Alternatively, the algorithm can be used to select nearest-neighbor parameters from a selection of modified bases and to calculate the stability of more than one sequence of the same length to give the same predetermined thermodynamic stability, with the selected modified bases.

Still further, the algorithm can be used to select one or more modified bases that allow the design of oligonucleotides of the same base pair length and substantially the same stability (T_m). These modified bases can be selected from a database containing predetermined thermodynamic nearest neighbor parameters from an collection of modified purines and/or modified pyrimidines. Preferably, the database contains nearest-neighbor parameters of 3-substituted pyrazolo[3,4-d]pyrimidines and/or 5-substituted pyrimidines.

Example 13

This example illustrates the use of modified oligonucleotides in an Invader™ assay.

5 The cleavase-based assay is shown in Figure 4 detects specific DNA and RNA sequences to cleave a complex formed by the hybridization of two overlapping oligonucleotides to a target. The enzyme cleaves the overhanging “flap”, which serves as an invader probe in the detection cassette probe where cleavage releases a fluorescence signal. The first cleavage takes places only when the single base mismatch in the invader is a perfect match. No reaction takes place with a mismatch target B. The cleaved “flap” serves as an invader in the detection cassette, leading to the release of fluorescence in the second cleavage step.

Table 7a and 7b

15 Comparison of Cleavase-bases amplification system with different modified bases substituted in the genomic and invader probes.

Table 7a

#	Probe ¹	Modified Bases Substituted in Probe Sequence ²	Length
1	Cassette	None	41
2	Invader T	None	59
3	Invader M1	Five A24 bases	32
4	Genomic T	None	43
5	Genomic M1	Six A14 bases	35
6	Genomic M2	Five A24 bases	31

¹ Unmodified Probes and assay conditions are similar to those described by Hall et al, *PNAS* 97:8272-77 (2000)

² A14 is hydroxypropynylPPA and A24 is 3-iododiaminoPPA

Table 7b

Genomic probe	Endpoint Match/Mismatch Signal Ratio	ΔF RFU ³
4	5	4100
5	10	8000
6	7	11900


³ ΔF is end point fluorescence difference between match and mismatch in relative fluorescence units

Table 7a shows a comparison of the effect of different modified bases when substituted in the Invader and genomic probes. The traditional invaderT (2) was substituted with 3-iododiaminoPPPA to give invader M1 which now has a length of 32 with a similar T_m as the 59-mer (2). Similarly the traditional genomic probe (4) was substituted 3-hydroxypropynylPPPA and 3-iododiaminoPPPA to give (5) and (6) respectively. Both of these probes are substantially shorter than the traditional (4). Using invaderM1 (3) in combination with the genomic probes 4-6 individually results in (Table 7b) improved performance of both genomic probes containing the modified bases (5 and 6) compared to the traditional genomic probe (4). As shown both the endpoint match/mismatch signal ratios and the end point fluorescence show increases compared to the unmodified genomic probe.


All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

WHAT IS CLAIMED IS:

- 1 1. A modified oligonucleotide comprising at least two bases selected
2 from the group consisting of unsubstituted and 3-substituted pyrazolo[3,4-d]pyrimidine
3 bases.
- 1 2. A modified oligonucleotide of claim 1, further comprising a covalently
2 attached minor groove binder.
- 1 3. A modified oligonucleotide of claim 1, further comprising at least one
2 covalently attached reporter group.
- 1 4. A modified oligonucleotide of claim 1, further comprising at least one
2 covalently attached quencher.
- 1 5. A modified oligonucleotide of claim 3, wherein said reporter group is a
2 fluorophore.
- 1 6. A modified oligonucleotide of claim 3, wherein said reporter group is a
2 fluorophore and said modified oligonucleotide further comprises an attached quencher.
- 1 7. A modified oligonucleotide of claim 1, comprising from 4 to 70 bases.
- 1 8. A modified oligonucleotide of claim 1, comprising from 4 to 70 bases
2 and further comprising an attached minor groove binder.
- 1 9. A modified oligonucleotide of claim 1, comprising from 4 to 70 bases
2 and further comprising an attached fluorophore and a quencher.
- 1 10. A modified oligonucleotide of claim 2, comprising from 4 to 70 bases
2 and further comprising an attached fluorophore and a quencher.
- 1 11. A modified oligonucleotide of claim 1, wherein at least one base of
2 said at least two bases is an unsubstituted pyrazolo[3,4-d]pyrimidine base having the formula:



4 and at least one of said at least two bases is a 3-substituted pyrazolo[3,4-d]pyrimidine base
5 having the formula:




6
7 wherein

8 each of said X¹ and X² groups is a member independently selected from the group
9 consisting of H, OH, NH₂ and a protected amino group; and

10 each of said R¹ groups is a member independently selected from the group consisting
11 of (C₁-C₁₂)heteroalkyl, (C₂-C₁₂)heteroalkenyl, (C₂-C₁₂)heteroalkynyl, -O-(C₁-
12 C₁₂)alkyl, -O-(C₂-C₁₂)alkenyl, -O-(C₂-C₁₂)alkynyl, -S-(C₁-C₁₂)alkyl, -S-(C₂-
13 C₁₂)alkenyl, -S-(C₂-C₁₂)alkynyl, heterocyclyl(C₁-C₁₂)alkyl, heterocyclyl(C₂-
14 C₁₂)alkenyl, heterocyclyl(C₂-C₁₂)alkynyl, aryl(C₁-C₁₂)alkyl, aryl(C₂-
15 C₁₂)alkenyl, aryl(C₂-C₁₂)alkynyl, aryl, heterocyclyl, halogen, -CN, -CONH₂
16 and protected forms thereof.

1 **12.** A modified oligonucleotide of claim 1, having the formula:



2
3 R² represents a first end of said modified oligonucleotide;

4 R³ represents a second end of said modified oligonucleotide;

5 the subscript n is an integer of from 4 to 70;


6 each B is a member independently selected from the group consisting of adenine,
7 thymine, cytosine, guanine, uracil, a pyrazolo[3,4-d]pyrimidine and a 3-
8 substituted pyrazolo[3,4-d]pyrimidine; and

9 each M is a member selected from the group consisting of an oligomer-forming sugar
10 and a peptide-nucleic acid-forming amino acid.


1 **13.** A modified oligonucleotide of claim 12, wherein at least one M is a
2 non-natural oligomer-forming sugar.

1 **14.** A modified oligonucleotide comprising at least one 5-substituted
2 pyrimidine base and at least one unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidine
3 base.

- 1 **15.** A modified oligonucleotide of claim **14**, further comprising a
2 covalently attached minor groove binder.
- 1 **16.** A modified oligonucleotide of claim **14**, further comprising at least one
2 covalently attached reporter group.
- 1 **17.** A modified oligonucleotide of claim **14**, further comprising at least one
2 covalently attached quencher.
- 1 **18.** A modified oligonucleotide of claim **16**, wherein said reporter group is
2 a fluorophore.
- 1 **19.** A modified oligonucleotide of claim **16**, wherein said reporter group is
2 a fluorophore and said modified oligonucleotide further comprises an attached quencher.
- 1 **20.** A modified oligonucleotide of claim **14**, comprising from 4 to 70
2 bases.
- 1 **21.** A modified oligonucleotide of claim **14**, comprising from 4 to 70 bases
2 and further comprising an attached minor groove binder.
- 1 **22.** A modified oligonucleotide of claim **14**, comprising from 4 to 70 bases
2 and further comprising an attached fluorophore and a quencher.
- 1 **23.** A modified oligonucleotide of claim **15**, comprising from 4 to 70 bases
2 and further comprising an attached fluorophore and a quencher.
- 1 **24.** A modified oligonucleotide of claim **14**, wherein said at least one 5-
2 substituted pyrimidine base having a formula selected from the group consisting of:




- 3
- 4 and said at least one unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidine base selected
5 from the group consisting of:



wherein

each of said X^1 , X^2 and X^3 groups is a member independently selected from the group consisting of H, OH, NH_2 and a protected amino group; and each of said R^1 and R^4 groups is a member independently selected from the group consisting of (C_1-C_{12}) heteroalkyl, (C_2-C_{12}) heteroalkenyl, (C_2-C_{12}) heteroalkynyl, $-O-(C_1-C_{12})$ alkyl, $-O-(C_2-C_{12})$ alkenyl, $-O-(C_2-C_{12})$ alkynyl, $-S-(C_1-C_{12})$ alkyl, $-S-(C_2-C_{12})$ alkenyl, $-S-(C_2-C_{12})$ alkynyl, heterocyclyl (C_1-C_{12}) alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl, aryl (C_2-C_{12}) alkenyl, aryl (C_2-C_{12}) alkynyl, aryl, heterocyclyl, halogen, $-CN$, $-CONH_2$ and protected forms thereof.

25. A modified oligonucleotide of claim 14, having the formula:



R^2 represents a first end of said modified oligonucleotide;

R^3 represents a second end of said modified oligonucleotide;


the subscript n is an integer of from 4 to 70;

each B is a member independently selected from the group consisting of adenine, thymine, cytosine, guanine, uracil, a 5-substituted pyrimidine and a 3-substituted pyrazolo[3,4-d]pyrimidine; and

each M is a member selected from the group consisting of an oligomer-forming sugar and a peptide-nucleic acid-forming amino acid.

26. A modified oligonucleotide of claim 25, wherein at least one M is a locked oligomer-forming sugar.

27. A modified oligonucleotide comprising from about 4 to about 70 bases and an attached minor groove binder, wherein at least one of said bases is replaced by a modified base selected from the group consisting of 5-substituted pyrimidines and unsubstituted or 3-substituted pyrazolo[3,4-d]pyrimidines.




wherein

each of said X^1 and X^2 groups is a member independently selected from the group consisting of H, OH, NH_2 and a protected amino group; and R^6 is a member selected from the group consisting of (C_1-C_{12}) heteroalkyl, (C_2-C_{12}) heteroalkenyl, (C_2-C_{12}) heteroalkynyl, $-O-(C_1-C_{12})$ alkyl, $-O-(C_2-C_{12})$ alkenyl, $-O-(C_2-C_{12})$ alkynyl, $-S-(C_1-C_{12})$ alkyl, $-S-(C_2-C_{12})$ alkenyl, $-S-(C_2-C_{12})$ alkynyl, heterocyclyl (C_1-C_{12}) alkyl, heterocyclyl (C_2-C_{12}) alkenyl, heterocyclyl (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl, aryl (C_2-C_{12}) alkenyl, aryl (C_2-C_{12}) alkynyl, aryl, heterocyclyl, halogen, $-CN$, $-CONH_2$ and protected forms thereof.

41. A modified oligonucleotide of claim **40**, wherein said heterocyclyl and aryl groups are selected from the group consisting of phenyl, tolyl, pyridyl, thiazolyl, imidazolyl, furanyl, oxazolyl, thienyl, pyrrolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indolyl, triazinyl, pyrimidinyl and naphthyl.


42. A modified oligonucleotide of claim **39**, wherein said minor groove binder has a formula selected from the group consisting of:



wherein

the subscript m is an integer of from 2 to 5;
the subscript r is an integer of from 2 to 10; and
each R^a and R^b is independently a linking group to said modified oligonucleotide, H, $-OR^c$, $-NR^cR^d$, $-COOR^c$ and $-CONR^cR^d$ wherein each R^c and R^d is selected from the group consisting of H, (C_1-C_{12}) heteroalkyl, (C_2-C_{12}) heteroalkenyl, (C_2-C_{12}) heteroalkynyl, (C_1-C_{12}) alkyl, (C_2-C_{12}) alkenyl, (C_2-C_{12}) alkynyl, aryl (C_1-C_{12}) alkyl and aryl.

43. A modified oligonucleotide of claim 41, wherein said minor groove binder has a formula selected from the group consisting of:




wherein

the subscript n is an integer of from 2 to 5;

the subscript r is an integer of from 2 to 10; and

each R^a and R^b is independently a linking group to said modified oligonucleotide, H, -OR^c, -NR^cR^d, -COOR^c and -CONR^cR^d wherein each R^c and R^d is selected from the group consisting of H, (C₁-C₁₂)heteroalkyl, (C₂-C₁₂)heteroalkenyl, (C₂-C₁₂)heteroalkynyl, (C₁-C₁₂)alkyl, (C₂-C₁₂)alkenyl, (C₂-C₁₂)alkynyl, aryl(C₁-C₁₂)alkyl and aryl.

44. A compound having the formula:



wherein


Z¹ is a member selected from the group consisting of H, F and OR^a wherein R^a is a member selected from the group consisting of H, (C₁-C₈)alkyl and a hydroxy protecting group;

Z² is a member selected from the group consisting of H and (C₁-C₈)alkyl, or optionally Z² is combined with Z¹ for form a five- to seven-membered ring;

Y¹ is a member selected from the group consisting of OH, a protected hydroxy group and O-P¹, wherein P¹ is a phosphoramidite or H-phosphonate group;

Y² is a member selected from the group consisting of OH, a protected hydroxy group and O-P², wherein P² is a phosphoramidite, H-phosphonate, monophosphate, diphosphate or triphosphate; and


B is a modified nucleotide selected from the group consisting of:




wherein

X^{11} and X^{12} are each independently selected from the group consisting of H, NH_2 and a protected amino group;
each R^{11} is independently selected from the group consisting of protected or unprotected forms of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl; and
each R^{12} is independently selected from the group consisting of protected or unprotected forms of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl, 3-methoxyprop-1-ynyl, 3-(hydroxymethyl)-4-hydroxy-1-butynyl, heterocyclyl(C_1 - C_{12})alkyl, heterocyclyl(C_2 - C_{12})alkenyl, heterocyclyl(C_2 - C_{12})alkynyl and heterocyclyl, with the proviso that R^{12} is other than 2-pyridylethynyl.


45. A compound of claim 44, wherein B is



46. A compound of claim 44, wherein B is




47. A compound of claim 44, wherein B is



2

1

48. A compound of claim **44**, wherein B is



2

1

49. A compound of claim **45**, wherein X¹¹ and X¹² are each NH₂.

1

50. A compound of claim **49**, wherein Y¹ is O-P¹, Y² is a protected hydroxy, Z¹ is H, R¹² is selected from the group consisting of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl.

2

3

1

51. A compound of claim **50**, wherein Y¹ is -O-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] and Y² is -O-(4,4'-dimethoxytrityl).

2

3

1

52. A compound of claim **45**, wherein X¹¹ is NH₂ and X¹² is H.

1

53. A compound of claim **52**, wherein Y¹ is O-P¹, Y² is a protected hydroxy, Z¹ is H, R¹² is selected from the group consisting of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl.

2

3

1

54. A compound of claim **53**, wherein Y¹ is -O-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] and Y² is -O-(4,4'-dimethoxytrityl).

2

3

1

55. A compound of claim **46**, wherein X¹² is H or NH₂.

1

56. A compound of claim **55**, wherein Y¹ is O-P¹, Y² is a protected hydroxy, Z¹ is H, R¹² is selected from the group consisting of 3-hydroxyprop-1-ynyl, 3-aminoprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl.

2

3

1

57. A compound of claim **56**, wherein X¹ is

2 -O-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] and X^2 is -O-(4,4'-
3 dimethoxytrityl).

1 58. A compound of claim 47, wherein X^{11} is NH_2 .


1 59. A compound of claim 58, wherein Y^1 is $O-P^1$, Y^2 is a protected
2 hydroxy, Z^1 is H, R^{11} is selected from the group consisting of 3-hydroxyprop-1-ynyl, 3-
3 aminoprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl.

1 60. A compound of claim 59, wherein Y^1 is
2 -O-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] and Y^2 is -O-(4,4'-
3 dimethoxytrityl).

1 61. A compound of claim 48, wherein Y^1 is $O-P^1$, Y^2 is a protected
2 hydroxy, Z^1 is H, and R^{11} is selected from the group consisting of 3-hydroxyprop-1-ynyl,
3 3-aminoprop-1-ynyl and 3-(hydroxymethyl)-4-hydroxy-1-butynyl.

1 62. A compound of claim 61, wherein Y^1 is
2 -O-[(2-cyanoethyl) N,N-diisopropylphosphoramidite] and Y^2 is -O-(4,4'-
3 dimethoxytrityl).

1 63. An oligonucleotide having the formula:



2
3 wherein


4 each Z^1 a member independently selected from the group consisting of H, F and
5 OR^a wherein R^a is a member selected from the group consisting of H, $(C_1-$
6 $C_8)$ alkyl and a hydroxy protecting group;

64. An oligonucleotide of claim **63**, wherein n is an integer of from 4

2 to 30.

1


65. An oligonucleotide of claim **63**, wherein at least one B is



2

1


66. An oligonucleotide of claim **63**, wherein at least one B is



2

1


67. An oligonucleotide of claim **63**, wherein at least one B is



2

1

68. An oligonucleotide of claim **63**, wherein at least one B is



2

1

69. An oligonucleotide of claim 63, wherein W¹ is a minor groove

2 binder-linking group moiety.

1

70. An oligonucleotide of claim **63**, wherein W¹ is a minor groove

2 binder-linking group moiety having an attached reporter group.

1

71. An oligonucleotide of claim 63, wherein W² is a minor groove

2 binder-linking group moiety.

(a) incubating a polynucleotide to be tested for the presence of the target sequence with a modified oligonucleotide having a sequence that is substantially complementary to the target sequence under hybridization conditions; and

(b) identifying hybridized nucleic acids;

wherein said modified oligonucleotide comprises at least one 3-substituted pyrazolo[3,4-*d*]pyrimidine in place of a purine residue.

90. A method in accordance with claim **89**, wherein said incubating is conducted in the presence of a cleavase enzyme.

91. A method in accordance with claim **89**, wherein said modified oligonucleotide further comprises a reporter group.

92. A method in accordance with claim **91**, wherein said reporter group is a fluorophore.


93. A method in accordance with claim **92**, said modified oligonucleotide further comprising an attached quencher.

94. A method in accordance with claim **89**, wherein said modified oligonucleotide further comprises an attached minor groove binder.


95. A method for primer extension, the method comprising incubating a polynucleotide containing a target sequence with one or more oligonucleotide primers complementary to the target sequence, in the presence of a polymerizing enzyme and nucleotide substrates under conditions favorable for polymerization; wherein at least one of the oligonucleotide primers contains a modified base selected from the group consisting of an unsubstituted pyrazolo[3,4-d]pyrimidine, a 3-substituted pyrazolo[3,4-*d*]pyrimidine and a 5-substituted pyrimidine base, in place of a purine or pyrimidine base.

96. A method in accordance with claim **95**, wherein one of said oligonucleotide primers is extended with a single base.


97. A method in accordance with claim 95, wherein said at least one of said oligonucleotide primers further comprises an attached minor groove binder.





PPPG




HOPPPG R = OH
NH₂PPPG R = NH₂




(NH₂)₂PPPA


$$\begin{aligned} &(\text{NH}_2)_2\text{PPPAOH} \quad \text{R} = \text{OH} \\ &(\text{NH}_2)_2\text{PPPANH}_2 \quad \text{R} = \text{NH}_2 \end{aligned}$$



PPPA



HOPPPA R = OH
NH₂PPPA R = NH₂



$\text{PU} \quad \text{R} = \text{H}$
 $\text{HOPU} \quad \text{R} = \text{OH}$



PC R = H
HOPC R = OH
NH₂PC R = NH₂

The wavy lines notes the position of an attached sugar or sugar derivative useful in oligonucleotide preparation.

Figure 2




Figure 2 is a graph illustrating the balancing of the T_m of 8-mer GC-rich and AT-rich probes with different combinations of MGB, A' and T' where A' is 3-propynylpyrazolopyrimidine and T' is 5-propynyluridine.

008277 6564260

5

10

15

20





Figure 3

Figure 4



00321 6567060

Variable	Mean	SD	Min	Max
Age	34.5	10.2	18	65
Gender	Male	100%		
Marital Status	Married	100%		
Education	High School	100%		
Occupation	Teacher	100%		
Income	\$15,000	\$5,000	\$10,000	\$25,000
Health Status	Good	100%		
Smoking Status	Non-smoker	100%		
Alcohol Consumption	None	100%		
Exercise Frequency	Weekly	100%		
Stress Level	Low	100%		
Sleep Quality	Good	100%		
Dietary Habits	Healthy	100%		
Family Size	2	1	1	4
Religious Beliefs	Christian	100%		
Political Views	Conservative	100%		
Travel Frequency	Monthly	100%		
Pet Ownership	Yes	100%		
Volunteering	Yes	100%		
Charitable Contributions	Yes	100%		
Community Involvement	Yes	100%		
Life Satisfaction	High	100%		
Overall Well-being	Excellent	100%		

Comparison of Invader Probs Performance with Different Number of PPGs

A. Six Gs substituted with PPG (tg*g*g*g*g*g*ctgtggcgctacg)

X-axis	6PPGs match	6PPGs mis
0	0.32	0.32
30	0.38	0.38
60	0.44	0.44
90	0.48	0.48
120	0.52	0.50
150	0.56	0.52
180	0.60	0.54
210	0.65	0.52


B. One G substituted with PPG (tgggg*ggcctggcggtacg)

X-axis	1PPG match	1PPG mis
0	0.32	0.32
30	0.36	0.36
60	0.40	0.40
90	0.44	0.44
120	0.48	0.48
150	0.52	0.50
180	0.56	0.52
210	0.62	0.48


B. No G substituted with PPG (tggggggcccctggcggtacg)

X-axis	all Gs match	all Gs mis
0	0.32	0.32
30	0.34	0.34
60	0.36	0.36
90	0.38	0.38
120	0.40	0.40
150	0.42	0.42
180	0.44	0.44
210	0.55	0.52


A. Six Gs substituted with PPG (tg*g*g*g*g*g*g*ccnnggcggctacg)



B. One G substituted with PPG (tgggg*ggccntggcggctacg)



B. No G substituted with PPG (tgggggggccugggcggcrag)



DECLARATION

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **MODIFIED OLIGONUCLEOTIDES FOR MISMATCH DISCRIMINATION** the specification of which X is attached hereto or was filed on as Application No. and was amended on (if applicable).

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application No.	Date of Filing	Status
09/640,953	08/16/2000	Pending
09/054,832	04/03/98	Pending
09/431,385	11/01/99	Pending
09/054,830	04/03/98	Patent
60/186,046	03/01/00	Pending

Full Name of Inventor 1:	Last Name: DEMPCY	First Name: ROBERT	Middle Name or Initial: O.	
Residence & Citizenship:	City: Kirkland	State/Foreign Country: Washington	Country of Citizenship: United States	
Post Office Address:	Post Office Address: 11421 NE 115th Court	City: Kirkland	State/Country: Washington	Postal Code: 98033
Full Name of Inventor 2:	Last Name: GALL	First Name: ALEXANDER	Middle Name or Initial: A.	
Residence & Citizenship:	City: Woodinville	State/Foreign Country: Washington	Country of Citizenship: Russia	
Post Office Address:	Post Office Address: 20051 170th Avenue, NE	City: Woodinville	State/Country: Washington	Postal Code: 98072
Full Name of Inventor 3:	Last Name: LOKHOV	First Name: SERGEY	Middle Name or Initial: G.	
Residence & Citizenship:	City: Kirkland	State/Foreign Country: Washington	Country of Citizenship: Russia	
Post Office Address:	Post Office Address: 13215 NE, 123rd Street, #312	City: Kirkland	State/Country: Washington	Postal Code: 98034
Full Name of Inventor 4:	Last Name: AFONINA	First Name: IRINA	Middle Name or Initial: A.	
Residence & Citizenship:	City: Mill Creek	State/Foreign Country: Washington	Country of Citizenship: United States	
Post Office Address:	Post Office Address: 2717 144th Court SE	City: Mill Creek	State/Country: Washington	Postal Code: 98012
Full Name of Inventor 5:	Last Name: SINGER	First Name: MICHAEL	Middle Name or Initial: J.	
Residence & Citizenship:	City: Seattle	State/Foreign Country: Washington	Country of Citizenship: United States	
Post Office Address:	Post Office Address: 11526 36th Avenue	City: Seattle	State/Country: Washington	Postal Code: 98125
Full Name of Inventor 6:	Last Name: KUTYAVIN	First Name: IGOR	Middle Name or Initial: V.	
Residence & Citizenship:	City: Woodinville	State/Foreign Country: Washington	Country of Citizenship: Russia	
Post Office Address:	Post Office Address: 18943 203rd Avenue, NE	City: Woodinville	State/Country: Washington	Postal Code: 98072
Full Name of Inventor 7:	Last Name: VERMEULEN	First Name: NICOLAAS	Middle Name or Initial: M.J.	
Residence & Citizenship:	City: Woodinville	State/Foreign Country: Washington	Country of Citizenship: United States	
Post Office Address:	Post Office Address: 19334 196th Avenue, NE	City: Woodinville	State/Country: Washington	Postal Code: 98072

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3
<u>Robert O. Dempcy</u>	<u>Alexander A. Gall</u>	<u>Sergey G. Lokhov</u>
Date	Date	Date
Signature of Inventor 4	Signature of Inventor 5	Signature of Inventor 6
<u>Irina A. Afonina</u>	<u>Michael J. Singer</u>	<u>Igor V. Kutyavin</u>
Date	Date	Date
Signature of Inventor 7		
<u>Nicolaas M.J. Vermeulen</u>		
Date		

SF 1162653 v1